

Jessica J. Chiang¹
 Shelley E. Taylor¹
 Julienne E. Bower^{1,2}

¹Department of Psychology
 University of California
 Los Angeles, CA
 E-mail: jchiang18@ucla.edu

²Cousins Center for
 Psychoneuroimmunology and Department
 of Psychiatry & Biobehavioral Sciences
 University of California
 Los Angeles, CA

Early Adversity, Neural Development, and Inflammation

ABSTRACT: Early adversity is a risk factor for poor mental and physical health. Although altered neural development is believed to be one pathway linking early adversity to psychopathology, it has rarely been considered a pathway linking early adversity to poor physical health. However, this is a viable pathway because the central nervous system is known to interact with the immune system via the hypothalamic-pituitary-adrenal (HPA) axis and autonomic nervous system (ANS). In support of this pathway, early adversity has been linked to changes in neural development (particularly of the amygdala, hippocampus, and prefrontal cortex), HPA axis and ANS dysregulation, and higher levels of inflammation. Inflammation, in turn, can be detrimental to physical health when prolonged. In this review, we present these studies and consider how altered neural development may be a pathway by which early adversity increases inflammation and thus risk for adverse physical health outcomes. © 2015 Wiley Periodicals, Inc. *Dev Psychobiol* 57:887–907, 2015.

Keywords: early adversity; amygdala; hippocampus; prefrontal cortex; hypothalamic-pituitary-adrenal axis; autonomic nervous system; inflammation

INTRODUCTION

A burgeoning literature suggests that early adverse experiences affect neural development. Compared to non-maltreated children, maltreated children show structural and functional differences in a number of neural regions, including the corpus callosum, amygdala, hippocampus, prefrontal cortex (PFC), and cerebellum, during both childhood and adulthood (Hart & Rubia, 2012; McCrory, De Brito, & Viding, 2012; Pechtel & Pizzagalli, 2011). Given that early adversity increases risk for developing deficits in cognitive

functioning and emotion processing and psychiatric disorders (Pechtel & Pizzagalli, 2011; Repetti, Taylor, & Seeman, 2002; Tyrka, Burgers, Philip, Price, & Carpenter, 2013), this area of research has focused on altered neural development as a potential pathway linking early adversity to poor mental health and psychopathology (Frodl & O'Keane, 2013; McCrory, De Brito, & Viding, 2011). However, altered neural development related to early adversity may also contribute to the link between early adversity and adverse physical health outcomes.

Psychosocial stress experienced in early life has a lasting effect on physical health. Adults who had cold, critical, abusive, and/or conflictual familial relationships in early life are more prone to develop obesity, lung disease, cancer, heart disease, and diabetes among other adverse physical health outcomes (Felitti et al., 1998; Miller, Chen, & Parker, 2011; Repetti et al., 2002). To better understand how early adversity comes to affect the development of poor physical health, researchers have focused their attention on underlying mechanisms. Elevated inflammation has emerged as a potential key pathway linking early adversity to poor physical health, as a number of studies have demonstrated a link between

Manuscript Received: 13 July 2014

Manuscript Accepted: 8 May 2015

Conflict of interest: none.

Correspondence to: Jessica J. Chiang

Contract grant sponsor: National Institute of Aging

Contract grant number: AG030309

Contract grant sponsor: National Science Foundation

Contract grant number: BCS-1124552

Contract grant sponsor: National Science Foundation Graduate Research Fellowship

Contract grant number: DGE-1144087

Article first published online in Wiley Online Library (wileyonlinelibrary.com): 15 September 2015

DOI 10.1002/dev.21329 • © 2015 Wiley Periodicals, Inc.

early adversity and elevated inflammation (Fagundes, Glaser, & Kiecolt-Glaser, 2013; Miller et al., 2011).

Inflammation is a natural, early response of the immune system to pathogens and injured tissue. In response to damaged tissue or infection, immune communication molecules known as pro-inflammatory cytokines, facilitate clearance of the pathogen and injured tissue from the body and promote repair. As such, inflammation is essential to healing and survival. However, prolonged inflammation can be harmful. For example, atherosclerosis is considered a disease rooted in inflammation, as pro-inflammatory cytokines are involved in its initiation and development (Libby, 2006; Pearson et al., 2003). Further, C-reactive protein (CRP), a marker of inflammation, is a robust and reliable predictor of cardiovascular disease with clinical cutoff points indicative of risk (Ridker, 2003). Inflammation has been implicated in a number of other chronic diseases as well, including depression (Raison, Capuron, & Miller, 2006), diabetes (Wellen & Hotamisligil, 2005), hypertension (Harrison et al., 2011), and certain cancers (Coussens & Werb, 2002; Hanahan & Weinberg, 2011; Rakoff-Nahoum, 2006).

The central nervous system and the immune system are known to interact, primarily via the hypothalamic-pituitary-adrenal (HPA) axis and autonomic nervous system (ANS) (Irwin & Cole, 2011; Rohleder, 2014). Thus, early adversity-related changes in neural development may have consequences for inflammatory processes and risk for poor health. Accordingly, the aim of this review is to detail how early adversity affects neural development in a manner that may be conducive to inflammation. A number of excellent reviews have synthesized the work on early adversity and neural development, with a focus on how adversity-related alterations in neural development and the HPA axis may increase risk for psychopathology (e.g., Danese & McEwen, 2012; Frodl & O'Keane, 2013; Hart & Rubia, 2012; Loman & Gunnar, 2010; Lupien, McEwen, Gunnar, & Heim, 2009; McCrory et al., 2012). There are also excellent reviews of research on early adversity and inflammation (Fagundes & Way, 2014; Miller et al., 2011), which tend to focus on physiological, molecular, psychosocial, and behavioral functioning as underlying mechanisms for adversity-related changes in inflammatory activity. The goal of this review is to integrate these two areas of work and specifically to highlight the role of neural development as a key pathway through which early adversity may influence downstream inflammatory processes. Thus, we build on past reviews by explicitly detailing the downstream effects of early adversity-related changes in neural development on systemic inflammation (Figure 1).

Although the multiple neurobiological systems interact with one another (as depicted in Figure 1), we focus on the effects of early adversity specifically on top-down regulation of inflammatory processes. More specifically, early adversity has been related to changes in neural regions that are part of a network that regulates HPA and ANS activity in the context of stress. The HPA axis and ANS, in turn, regulate inflammatory processes. As we outline below, altered neural development in these regions can lead to more frequent, prolonged, and/or exaggerated HPA and ANS responses to subsequent psychosocial stress. Over time, such responses can lead to the dysregulation of the HPA axis and ANS. Because the HPA axis and ANS regulate inflammatory processes, their dysregulation over prolonged periods of time can promote inflammation, thereby increasing risk for poor physical health outcomes.

We present evidence for this pathway by first providing an overview of the connection between the central nervous and inflammatory systems, focusing on the amygdala, hippocampus, and PFC. We focus on these regions because they have been widely studied and have been established as key neural regions implicated in stress processes, although other neural regions (e.g., periaqueductal gray of midbrain, hypothalamus) are also implicated in stress processes. Next, we selectively review studies supporting various pieces of the pathway, namely those that have linked early adversity to neural development, HPA axis and ANS functioning, and inflammation. We then integrate these findings and consider how altered neural development might ultimately influence inflammatory processes. We also note the reciprocal relations between neural development and inflammatory processes by briefly describing the role of inflammation on neural development, although this topic is beyond the scope of this article (for reviews on this topic, see Deverman & Patterson, 2009; Musaeelyan et al., 2014; Stolp, 2013). Other related topics that are beyond the scope of the present review include glucocorticoids as a mechanism of the effects of early adversity on neural development (Lupien et al., 2009; Tottenham & Sheridan, 2010), and early attachment learning involving the HPA axis and amygdala (Moriceau, Roth, & Sullivan, 2010). We conclude by raising several issues that future research should address in order to advance our understanding of the detrimental effects of early adversity.

Although children can experience a wide range of stressors, such as socioeconomic disadvantage and natural disasters, we primarily draw on studies examining child maltreatment (i.e., abuse and neglect) and other aspects of the family environment, including




FIGURE 1 Neurodevelopmental pathway by which early adversity may result in elevated inflammation.

family conflict and poor maternal caregiving, given that much of the extant literature on the effects of early life stress on neural development has focused on these types of stressors. Importantly, because these aspects of the family environment are typically enduring and may have long-term effects on physiology (Juster, McEwen, & Lupien, 2010), we focus on early life stress that is chronic. Although we use child maltreatment as the representative construct for understanding the effects of early adversity, our use of “early life stress” and “early adversity” is not confined to child maltreatment, but rather, is used as a broader term that can encompass other chronic psychosocial stressors.

NEURAL REGULATION OF INFLAMMATION

The central nervous system influences inflammatory processes through its regulation of the HPA axis and ANS. Perception of threat initiates the experience of stress, which triggers activation of the HPA axis and sympathetic nervous system (SNS) and withdrawal of the parasympathetic nervous system (PNS). The amygdala, hippocampus, and PFC are key components of the integrated neural network that matches appraisal and interpretation of potentially threatening environmental stimuli to behavioral and biological responses to threat (Fuchs & Flügge, 2003). As such, these neural regions regulate ANS and HPA activity. The HPA axis and ANS, in turn, regulate inflammatory processes.

Central Regulation of the HPA Axis and ANS

The amygdala and hippocampus are particularly important for detecting both physical and psychological threat (e.g., Faselow & Gale, 2003; LeDoux, 2003). The amygdala plays a critical role in fear-conditioning, which involves pairing novel and initially neutral stimuli with an emotional response. Through such classical conditioning, one learns which aspects of his or her environment are dangerous. Thus, perception of threat activates the amygdala. When threat is perceived, the amygdala triggers the HPA axis and ANS through its projections to the hypothalamus and the brainstem (Gianaros & Sheu, 2009; Herman et al., 2003).

Appraisal of novel, potentially threatening situations is partially a function of learning and memories of past events, including actual aversive experiences and vicarious experiences shared through verbal communication with others (Phelps, 2004). Involved in learning, contextual fear conditioning (Rudy, 1993) and memory (Chadwick, Hassabis, Weiskopf, & Maguire, 2010; Squire, 1992), the hippocampus is crucial for recognizing threat in the environment. Furthermore, the hippocampus interacts with the amygdala to facilitate retrieval of emotionally valenced contextual information (Smith, Stephan, Rugg, & Dolan, 2006), which plays a role in appraisal and reactivity to subsequent contextual stimuli (Maren, Phan, & Liberzon, 2013). As such, the hippocampus can indirectly activate the HPA axis and ANS by modulating amygdala activation (Phelps, 2004). In addition, the hippocampus is directly

involved in inhibiting the HPA axis (Herman et al., 2003), as described in more detail below.

The medial and ventral PFC regions are involved in fear extinction, whereby responses to previously acquired fear dissipate after repeated exposure to a conditioned stimulus in the absence of the unconditioned stimulus (Sotres-Bayon, Cain, & LeDoux, 2006). Thus, medial and ventral PFC activation downregulates amygdala activity, inhibiting responses to acquired fear and other threats (Hariri, Mattay, Tessitore, Fera, & Weinberger, 2003; Quirk, Likhtik, Pelletier, & Paré, 2003; Sotres-Bayon et al., 2006). The medial PFC also connects to and interacts with the hippocampus during retrieval of extinction memory (Maren et al., 2013; Milad et al., 2007; Orsini, Kim, Knapska, & Maren, 2011), an important aspect of appraisal processes. Given amygdalar and hippocampal involvement in threat perception and initiation of HPA and ANS responses to threat, ventral and medial PFC regulation of the amygdala and interactions with the hippocampus may attenuate HPA and ANS responses to stress and/or facilitate physiological recovery from stress. Rich in glucocorticoid receptors, the PFC also plays a role in suppressing the HPA axis (Diorio, Viau, & Meaney, 1993; Sullivan & Gratton, 2002). Interestingly, the PFC has also been shown to exert excitatory control over the HPA axis (Radley, Arias, & Sawchenko, 2006; Wang et al., 2005). Whether the PFC excites or inhibits the HPA axis may depend on the particular area of the PFC and type of stressor (Herman et al., 2003; Herman & Mueller, 2006; Kern et al., 2008; Sullivan & Gratton, 2002).

HPA Axis and ANS Regulation of Inflammation

Neural processing of stress leads to changes in the HPA axis and ANS. More specifically, perception of threat activates the amygdala, which signals the hypothalamus to secrete corticotrophin releasing hormone (CRH). CRH stimulates the pituitary to secrete adrenocorticotrophic hormone, which then stimulates the adrenal cortex to produce and release cortisol. In a negative feedback loop, cortisol binds to glucocorticoid receptors in several brain regions, including the hippocampus, PFC, amygdala, and hypothalamus, suppressing CRH production and the HPA stress response. When bound to glucocorticoid receptors on immune cells, cortisol can suppress transcription of pro-inflammatory genes and subsequent production and release of proinflammatory cytokines (Sternberg, Chrousos, Wilder, & Gold, 1992). Cortisol, then, is anti-inflammatory and protects against negative effects of an exaggerated or sustained inflammatory response (Glezer & Rivest, 2004; Irwin & Cole, 2011; Rohleder, 2011). Indeed, low levels of cortisol

have been related to higher systemic levels of inflammatory markers (Kunz-Ebrecht, Mohamed-Ali, Feldman, Kirschbaum, & Steptoe, 2003; Rohleder, 2014). Importantly, in the context of stress, sustained activation of the HPA axis can desensitize glucocorticoid receptors (Avitsur, Stark, & Sheridan, 2001; Cohen et al., 2012; Miller, Cohen, & Ritchey, 2002; Sheridan, Stark, Avitsur, & Padgett, 2000). Consequently, inflammatory cytokines may not “hear” the signal of cortisol, failing to respond to cortisol’s inhibitory signals (Miller et al., 2002; Rohleder, 2011). Desensitized glucocorticoid receptors, then, may also lead to inadequate suppression of inflammation.

Perception of threat also activates the SNS and triggers withdrawal of the PNS. Activation of the amygdala stimulates the neurons in the paraventricular nucleus of the hypothalamus, which projects to sympathetic preganglionic cells in the thoracic spinal cord (Zagon & Smith, 1993) both directly and indirectly via connections to the brainstem. Axons of these cells synapse with those of the adrenal medulla, which secretes the catecholamines, epinephrine and norepinephrine, into systemic circulation. Perception of stress also releases norepinephrine into local microenvironments of vasculature, organs, and tissues via SNS neural fiber innervation (Irwin & Cole, 2011). Norepinephrine, in turn, activates adrenergic receptors. Although the anti- or pro-inflammatory effect of norepinephrine depends on a number of factors, including the adrenergic receptor subtype that it activates, specific tissue, and local concentration of norepinephrine (McNamee et al., 2010; Yang, Lee, Kim, Suh, & Chong, 2012), it can upregulate pro-inflammatory gene expression in monocytes (Cole et al., 2010; Grisanti et al., 2011; Irwin & Cole, 2011). In particular, when bound to α_2 and β adrenergic receptors, norepinephrine stimulates NF- κ B, a key transcriptional pathway that increases inflammation (Bierhaus et al., 2003). Thus, activation of the SNS in response to stress can increase inflammation.

The paraventricular nucleus also projects directly, and indirectly via connections to the brainstem, to preganglionic neurons of the vagus nerve, the primary nerve of the PNS. Activation of the PNS releases the vagal neurotransmitter acetylcholine. When bound to α_7 nicotinic receptors, acetylcholine suppresses pro-inflammatory cytokine production (Abboud, Harwani, & Chappelle, 2012; Rohleder, 2011; Thayer, 2009; Tracey, 2002, 2009). Under tonic conditions, the PNS exerts inhibitory control (Thayer, Ahs, Fredrikson, Sollers III, & Wager, 2012). In the context of stress, parasympathetic activity is typically diminished (Mueller et al., 2012; Thayer, Yamamoto, & Brosschot, 2010; Weber et al., 2010). Thus, low vagal tone and decreased

vagal withdrawal during psychological stress have been associated with increased inflammation (e.g., Haensel, Mills, Nelesen, Ziegler, & Dimsdale, 2008; Thayer & Fischer, 2009; Weber et al., 2010).

In sum, the amygdala, hippocampus, and PFC are central components of the neural network underlying stress processes. Consequently, they can influence inflammatory processes through the HPA axis and ANS. The amygdala and hippocampus are crucial to perceiving and recognizing threat in the environment and can therefore activate biological stress systems. The PFC inhibits biological stress responses by down-regulating amygdala activity. In addition, the hippocampus and PFC play a role in the negative feedback loop of the HPA axis. The HPA axis and ANS can impact inflammatory processes, with the HPA axis and PNS typically exerting anti-inflammatory and the SNS exerting pro-inflammatory effects.

EARLY ADVERSITY AND NEURAL DEVELOPMENT

Experience customizes neural connections and their interactions via synaptic formation and pruning of excess neural connections, a mechanism thought to maximize chances of survival in a given environment (Fox, Levitt, & Nelson III, 2010; Greenough, Black, & Wallace, 1987). Although neural systems and processes are sensitive to experiences throughout the lifespan (Pascual-Leone, Amedi, Fregni, & Merabet, 2005), they may be particularly sensitive to experiences early in life when they undergo significant and rapid development (Lupien et al., 2009). The amygdala, hippocampus, and PFC undergo dramatic changes during childhood and adolescence (Blakemore & Choudhury, 2006; Tottenham & Sheridan, 2010). Therefore, adverse experiences during these periods have been shown to have long-lasting effects on these stress-related neural regions.

Functional Changes

A number of studies have related early adversity to functional changes in the amygdala, hippocampus, and PFC both in childhood and adulthood. Compared to children and adolescents who were never institutionalized, previously institutionalized children and adolescents show greater amygdala responses to threatening interpersonal cues (i.e., fearful and angry faces) (Gee et al., 2013; Maheu et al., 2010; Tottenham et al., 2011). Initial evidence suggests that early adversity may also be related to altered hippocampal and PFC functioning. Specifically, previously institutionalized

children have exhibited greater hippocampal reactivity to psychosocial threat (Maheu et al., 2010) and lower glucose metabolism in the hippocampus and PFC (Chugani et al., 2001). Behavioral studies have also shown that early institutionalization is associated with deficits in hippocampal-dependent memory (Bos, Fox, Zeanah, & Nelson III, 2009). Notably, the previously institutionalized children and adolescents of these studies were subsequently adopted, which suggests that the effects of early adversity on amygdala, hippocampal, and PFC functioning persist despite termination of the stressor (Maheu et al., 2010; Tottenham et al., 2011). With the exception of the studies just described, relatively few studies have examined how early adversity impacts PFC and hippocampal functioning in children; more evidence exists for the link between early adversity and altered amygdala functioning in children. Further, these studies have focused on previously institutionalized youth, which represents a relatively extreme case of neglect and caregiver deprivation. Whether these findings generalize to youth experiencing other types of adversity remains unclear. Nevertheless, these studies collectively begin to delineate how early adversity impacts amygdala, hippocampal, and PFC functioning in youth.

These disruptions in neural functioning appear to persist into adulthood. For instance, abuse, neglect, and a cold, unsupportive childhood family environment have been associated with greater amygdala reactivity to interpersonal threat-related cues in adults (Dannlowski et al., 2013; Dannlowski et al., 2012; Taylor, Eisenberger, Saxbe, Lehman, & Lieberman, 2006; van Harmelen et al., 2012). Although research on how early adversity may alter hippocampal and PFC functioning in adulthood is scant, preliminary evidence suggests that these relations exist. In one study of 18 adults, childhood maltreatment was associated with greater hippocampal responses to novel, potentially threatening interpersonal cues in adulthood (Edmiston & Blackford, 2013). Behavioral studies offer more support for this link. A history of sexual abuse and greater number of various childhood adverse experiences have been associated with deficits in hippocampal-dependent learning and memory, including short-term, verbal, visual, autobiographical, and global memory scores, slower learning, and reduced persistence during extinction after partial reinforcement (Anda et al., 2006; Navalta, Polcari, Webster, Boghossian, & Teicher, 2006; Patterson, Craske, & Knowlton, 2013). In another study, a cold, harsh, conflict-ridden family environment has been associated with greater right ventrolateral PFC responses to threat-related cues, but this greater PFC activity was correlated with greater rather than decreased amygdala activity, indicating a deficit in PFC

regulation of amygdala responses to threat (Taylor et al., 2006). Together, these studies indicate that changes in amygdala, hippocampal, and PFC functioning resulting from early adversity may be enduring.

Structural Changes

Prior research has also demonstrated an association between early adversity and changes in neural structure during both childhood and adulthood. Neglected and abused children and adolescents have been shown to have greater amygdala volume (Mehta et al., 2009; Tottenham et al., 2010) and smaller anterior cingulate cortex and orbital frontal cortex volume, both subregions of the PFC (De Brito et al., 2013; Hanson et al., 2010; Kelly et al., 2013), compared to children who have not been maltreated. Similarly, irrational punishment, high levels of family conflict, and maternal depressive symptomatology (which is associated with poor maternal care), and cumulative life stress have been associated with larger amygdala and smaller PFC volume in children and adolescents (Edmiston et al., 2011; Hanson et al., 2012; Korgaonkar et al., 2013; Lupien et al., 2011; Whittle et al., 2009). Interestingly, early adversity has not been related to structural changes in the hippocampus during childhood and adolescence (Frodl & O'Keane, 2013; Lupien et al., 2011).

In adulthood, early adversity is related to decreased hippocampal and PFC volume, but not to amygdala volume (Baker et al., 2013; Cohen et al., 2006; Korgaonkar et al., 2013). Healthy adults who experienced neglect and/or abuse in childhood had smaller hippocampal white and gray matter (Chaney et al., 2014; Dannlowski et al., 2012; Frodl, Reinhold, Koutsouleris, Reiser, & Meisenzahl, 2010) and decreased mPFC gray matter (Dannlowski et al., 2012; van Harmelen et al., 2010), compared to their non-maltreated counterparts. Other types of early life stressors including living with a mentally ill family member, exposure to violence, harsh discipline, and parental separation have also been associated with decreased hippocampal and PFC volume in adulthood (Teicher, Anderson, & Polcari, 2012) and PFC volume (Andersen et al., 2008; Cohen et al., 2006; Frodl et al., 2010; Kitayama, Quinn, & Bremner, 2006; Tomoda et al., 2009; Teicher, Anderson, & Polcari, 2012; Treadway et al., 2009).

Overall, evidence suggests that early adversity can lead to persistent functional changes in the amygdala, hippocampus, and PFC. Early adversity also appears to alter neural structure, though some of these effects may be observable only during particular periods of the lifespan. Specifically, changes in amygdala volume as a

result of early adversity are apparent earlier in life, but do not seem to persist into adulthood. This may be because early adverse experience precociously engages the amygdala and accelerates its development (for review, see Roozendaal, McEwen, & Chattarji, 2009; Tottenham & Sheridan, 2010). By contrast, the link between early life stress and alterations in hippocampal structural development is not apparent in childhood and adolescence, but seems to emerge in adulthood. Although the mechanisms have yet to be elucidated, it is possible that any effects on pruning of synapses that may lead to decreased hippocampal volume may not be observable until later in life (Andersen & Teicher, 2004, 2009).

Curiously, very few studies have examined the effects of early adversity on the connectivity and interactions of the amygdala, hippocampus, and PFC despite that they collectively regulate responses to stress. An exception is a recent study (Herringa et al., 2013) that found that previous maltreatment was associated with decreased hippocampal-mPFC and amygdala-mPFC connectivity. Although this study offers initial support for the negative effects of early adversity on neural connectivity, only female adolescents were examined. Whether early adversity impacts neural connectivity in children and persists into adulthood is currently unknown.

EARLY ADVERSITY AND HPA AND ANS DYSREGULATION

The preceding section described evidence for the link between early adversity and alterations in the amygdala, hippocampus, and PFC. Given that these regions play an essential role in regulating the HPA axis and ANS, altered development in these regions may lead to altered HPA axis and ANS activity. We would expect, then, that early adversity be related to altered HPA axis and ANS functioning. In this section, we present findings documenting links between early adversity and HPA and ANS dysregulation.

HPA Axis

Because other reviews (e.g., Gunnar & Quevedo, 2007; Repetti, Robles, & Reynolds, 2011) have addressed the effects of early adversity on the HPA axis in detail, we briefly summarize findings. There is compelling evidence that early adversity leads to alterations in HPA functioning in childhood, adolescence, and adulthood. However, the direction of the effect is complex, as studies have documented both hypoactive and hyperactive HPA activity following early adversity. For

instance, children with depressed mothers or who were maltreated have lower basal levels of cortisol and blunted cortisol responses to challenge (Dietz et al., 2013; Fernald, Burke, & Gunnar, 2008; Ouellet-Morin et al., 2011). This blunted pattern of cortisol secretion in response to acute stress is also evident among adolescents and adults who have experienced early life stress (e.g., Carpenter et al., 2007; Carpenter, Shattuck, Tyrka, Geraciotti, & Price, 2011; Carpenter et al., 2009; Engert et al., 2010b; Lovallo, Farag, Sorocco, Cohoon, & Vincent, 2012; Luecken, Kraft, & Hagan, 2009; MacMillan et al., 2009; Taylor, Karlamangla, Friedman, & Seeman, 2011; Trickett, Gordis, Peckins, & Susman, 2014).

Paradoxically, early adversity has also been linked to exaggerated levels of cortisol secretion. Specifically, neglect and maternal stress and depression, which increase maternal neglect, hostility, and physical assault (Lovejoy, Graczyk, O'Hare, & Neuman, 2000; Turney, 2011), have been associated with higher basal cortisol and cortisol responses to an acute stressor in children (Ashman, Dawson, Panagiotides, Yamada, & Wilkinson, 2002; Essex, Klein, Cho, & Kalin, 2002; Sullivan, Bennet, & Lewis, 2013). High levels of basal and stress-induced cortisol as a result of early adversity are also evident in adulthood (e.g., Engert et al., 2010a; Heim et al., 2000, 2002). For example, maternal separation in childhood has been related to a larger cortisol awakening response and a flatter diurnal cortisol slope (Kumari, Head, Bartley, Stansfeld, & Kivimaki, 2012).

Together, these studies show that early adversity can have enduring effects on HPA function, although research has yet to disentangle factors that lead to different trajectories of HPA functioning. Variation in severity and type of stressors and in methods for assessing HPA functioning across studies may help explain the divergent findings (Dietz et al., 2013; Doom, Cicchetti, & Rogosch, 2014; Struber, Struber, & Rother, 2014). For instance, one study found that adults who were severely maltreated had lower morning cortisol levels whereas moderately maltreated adults had higher cortisol levels (van der Vegt, van der Ende, Kirschbaum, Verhulst, & Tiemeier, 2009). In another study, physical abuse was related to quicker cortisol reactivity to stress whereas emotional abuse was related with delayed cortisol recovery from stress (Kuhlman, Geiss, Vargas, & Lopez-Duran, 2015). Timing of stress and assessment of HPA function may also contribute to the mixed findings, as age and pubertal development can affect HPA function (Gunnar, Wewerka, Frenn, Long, & Griggs, 2009; Romeo, 2010; Stroud et al., 2009). Studies of early adversity that repeatedly assess HPA function through childhood into adulthood can

help clarify the effects of early adversity on HPA hyper- and hypo-secretion. Importantly, both exaggerated and blunted cortisol levels generally indicate HPA dysregulation (McEwen, 1998). HPA dysregulation, in turn, can lead to glucocorticoid receptor insensitivity (Miller et al., 2002), a state that may ultimately determine cortisol's anti-inflammatory effects (Cohen et al., 2012). Preliminary evidence suggests that early adversity desensitizes glucocorticoid receptors (Miller & Chen, 2010; Miller et al., 2009), which could explain a state of increased inflammation despite high levels of cortisol among some individuals with a history of early adversity.

ANS

Fewer studies have examined the effects of early adversity on ANS functioning. Nevertheless, those studies suggest that early adversity may affect sympathetic functioning beginning in childhood and persisting into adulthood. Specifically, cross-sectional and longitudinal studies have shown that children exposed to neglect, family stress (e.g., parental depression, anger, and role overload, and financial and parenting stress), and aggressive marital conflict exhibit greater sympathetic activity to laboratory challenges (El-Sheikh, 2005; Ellis, Essex, & Boyce, 2005; Oosterman, De Schipper, Fisher, Dozier, & Schuengel, 2010). As adults, individuals who experienced stress (i.e., abuse, neglect, high levels of family conflict, illness, loss) in early life exhibit increases in heart rate and blood pressure both at rest and in response to acute stress (Gatt et al., 2009; Heim et al., 2000; Larkin, Frazer, & Semenchuk, 1996; Larkin, Frazer, & Wheat, 2011; Lovallo, 2013; Luecken, 1998; Taylor, Lerner, Sage, Lehman, & Seeman, 2004), although not all studies have not found this effect (Larkin et al., 2011; Luecken & Roubinov, 2012).

Early adversity also seems to have an impact on parasympathetic functioning. Marital conflict and negative, hostile, controlling, and cold maternal behavior have been associated with low vagal tone (i.e., the continuous PNS activity under tonic conditions) and decreased vagal withdrawal (i.e., diminished PNS activity under challenging conditions), in infants and children (Calkins, Graziano, Berdan, Keane, & Degnan, 2008; Calkins, Smith, Gill, & Johnson, 1998; Katz, 2007; Moore, 2010; Porter, Wouden-Miller, Silva, & Porter, 2003). The majority of these studies are cross-sectional and examine children age six or younger. The relation between early adversity and parasympathetic functioning in later childhood, adolescence, and adulthood has been less examined. However, initial evidence suggests that early adversity may have a long-lasting

effect on parasympathetic functioning. Two studies of female adolescents found that documented history of maltreatment prospectively predicts lower vagal tone (Miskovic, Schmidt, Georgiades, Boyle, & MacMillan, 2009; Shenk, Noll, Putnam, & Trickett, 2010). Similarly, one study (Dale et al., 2009) found that adult women with a history of abuse exhibited poorer vagal regulation during and immediately after physical stress. The modest evidence necessitates additional investigation to conclude whether early adversity has effects on PNS functioning beyond early childhood and if so, whether it has similar effects on the PNS in males.

In sum, some evidence suggests that early adversity is related to exaggerated SNS activity and poor vagal regulation both in childhood and adulthood. However, such studies are relatively few, and more research is needed to confirm that these associations persist into adulthood. One of the complications in this area of research is the focus on heart rate and blood pressure, which are influenced by both the sympathetic and parasympathetic nervous systems. Studies that use “pure” measures of SNS activity (e.g., pre-ejection period) may clarify the enduring impact of early life stress on this system. Stronger support exists for persistent HPA dysregulation following early adversity, as numerous studies have documented aberrant cortisol levels related to early adversity in youth and adults. Though both hyper and hypo-cortisolism may be indicators of HPA dysregulation, perhaps more important for inflammation is the sensitivity of glucocorticoid receptors, and emerging evidence suggests that early adversity may also render glucocorticoid receptors insensitive.

EARLY ADVERSITY AND INFLAMMATION

The HPA axis and ANS modulate inflammatory processes, as described above. HPA axis and ANS dysregulation resulting from early adversity, then, can affect inflammatory activity. Consequently, early adversity should be related to altered inflammatory activity. In this section, we summarize evidence for this notion.

A growing number of studies suggest that throughout the lifespan, individuals maltreated in childhood experience greater inflammatory activity, both tonically and in response to psychosocial stress, in the absence of microbial threat and disease. For example, in one prospective study (Slopen, Kubzansky, McLaughlin, & Koenen, 2013), mothers reported whether their children experienced any socially adverse events (i.e., taken into foster care, physically hurt by someone, sexually abused, separated from either parent) in the previous 12–18 months every year for the first eight years of their children’s lives. Children’s levels of circulating

interleukin-6, a pro-inflammatory molecule, and CRP, were then assessed at ages 10 and 15. Cumulative adverse events were associated with greater levels of inflammation in both childhood and adolescence. Similarly, childhood bullying was prospectively related to higher levels of circulating levels of CRP in childhood and adolescence (Copeland et al., 2014). Such prospective studies of early adversity and inflammation are rare, and some studies have not found a link between early adversity and circulating markers of inflammation in plasma (e.g., Miller & Chen, 2010). Nevertheless, the longitudinal studies (Copeland et al., 2014; Slopen et al., 2013) described offer strong support for the link between early adversity and inflammation in youth given that they employed a prospective design and included large samples ($n > 1000$) and repeated assessments of adversity throughout childhood.

Studies of adults with a history of early adversity have more consistently shown that early adversity is associated with heightened inflammation. As adults, people who were raised in a cold, conflictual family environment (e.g., harsh discipline, maternal rejection) or experienced child maltreatment or caregiver changes have greater circulating levels of CRP, as demonstrated cross-sectionally (Kiecolt-Glaser et al., 2011; Schrepf, Markon, & Lutgendorf, 2014; Taylor, Lehman, Kiefe, & Seeman, 2006). When challenged, adults with a history of early adversity exhibit an exaggerated inflammatory response. Specifically, adults who experienced childhood maltreatment compared to those who did not had greater increases in circulating concentrations of interleukin-6 in response to a laboratory social stressor (Carpenter et al., 2010; Pace et al., 2006) and to daily stress (Gouin, Glaser, Malarkey, Beversdorf, & Kiecolt-Glaser, 2012). The majority of these studies rely on retrospective reports of early life experiences, raising the concern of recall biases. However, prospective studies with repeated assessments of early adversity using parental reports, child reports, and behavioral observations have also found that early adversity is related to greater circulating levels of inflammatory markers in adulthood (Copeland et al., 2014; Danese, Pariante, Caspi, Taylor, & Poulton, 2007; Lacey, Kumari, & Bartley, 2014; Matthews, Chang, Thurston, & Bromberger, 2014). These studies offer compelling evidence that early adversity is related to heightened inflammation in adulthood.

FROM ALTERED NEURAL DEVELOPMENT TO INCREASED INFLAMMATION

Thus far, we have described how the central nervous system links to inflammatory processes and have

reviewed past studies showing associations between early adversity and altered neural development, HPA axis and ANS dysregulation, and increased inflammation. But exactly how might adversity-related alterations in neural development increase inflammatory activity? Based on the lines of research reviewed above, we consider the downstream effects of altered neural development on systemic inflammation in this section.

As described above, the amygdala and hippocampus are involved in detecting and recognizing threat in the environment. Thus, a hypersensitive amygdala and deficits in hippocampal-dependent learning and memory processes resulting from early life stress can lead to more frequent detection of threat in the environment. Behavioral studies have supported this notion, showing that children who were maltreated or perceived less warmth and acceptance from their parents appraised their environment as more threatening (Davies & Cicchetti, 2004; Pollak & Kistler, 2002) and perceived more social stress in their adolescent years (Raposa, Hammen, Brennan, O'Callaghan, & Najman, 2014).

Because perception of threat activates physiological stress systems (Gaab, Rohleder, Nater, & Ehlert, 2005), the functional alterations in the amygdala and hippocampus may be accompanied by repeated, exaggerated, and/or prolonged activation of the biological stress systems. Moreover, given that the mPFC downregulates amygdala activity, early adversity that disrupts mPFC function and its connectivity to the amygdala could potentially lead to inadequate inhibition of amygdala activity. This would further sustain prolonged and/or exaggerated responses to threat. Consistent with this notion, greater amygdala activity has been related to greater HPA axis and SNS activity, and to lesser PNS activity (Muscatell & Eisenberger, 2012; Thayer et al., 2012; Urry et al., 2006). By contrast, increased activity in the mPFC has been related to lower levels of cortisol (Kern et al., 2008), decreased blood pressure and heart rate (Hänsel & von Känel, 2008; Kern et al., 2008; Pruessner et al., 2008; Verberne & Owens, 1998; Wager et al., 2009), and increased vagal activity (Lane et al., 2009; Thayer et al., 2012). Moreover, a stronger coupling of the medial PFC and the amygdala is predictive of a steeper decline in the daily rhythm of cortisol (Urry et al., 2006), which is believed to be more adaptive.

Structural changes in the brain may also alter HPA function, although the mechanisms are not entirely clear. Support for this notion comes primarily from studies on animals and on patients sustaining brain damage or lesions. Hippocampal lesions or damage in rodents and humans results in elevated basal levels of glucocorticoids (Fendler, Karmos, & Telegdy, 1961; Knigge, 1961), disrupts the diurnal rhythm of glucocor-

ticoids (Buchanan, Kern, Allen, Tranel, & Kirschbaum, 2004; Fischette, Komisaruk, Edinger, Feder, & Siegel, 1980; Herman & Mueller, 2006), and delays HPA recovery after stress (Herman, Cullinan, Morano, Akil, & Watson, 1995; Magarinos, Somoza, & De Nicola, 1987; Sapolsky, Armanini, Sutton, & Plotsky, 1989). Human studies of non-clinical samples have also shown that smaller hippocampal volume is related to a disrupted diurnal pattern of cortisol (Buchanan et al., 2004; Pruessner et al., 2010). Similarly, animal studies show that damage or lesions to the mPFC can result in an enhanced HPA response to acute stress (Brake et al., 2000; Diorio et al., 1993; Figueiredo, Bruestle, Bodie, Dolgas, & Herman, 2003; Herman & Cullinan, 1997). Decreased PFC volume has also been associated with a failure to suppress cortisol responses to dexamethasone challenge (MacLulich et al., 2006), and higher levels of cortisol at bedtime when cortisol levels should be approaching their nadir (Carrion, Weems, Richert, Hoffman, & Reiss, 2010). Taken together, these studies show that decreased hippocampal and PFC volume can disrupt HPA axis functioning. Whether alterations in hippocampal and PFC structure affects ANS function is currently unclear. Nevertheless, to the extent that neural structure underlies neural function, smaller hippocampal and PFC volume may also affect ANS activity.

Given that the HPA axis and ANS regulate inflammatory processes, we would expect that stress-related changes in neural function and structure would relate to inflammation. To date, only two human studies to our knowledge have examined the link between neural activity during stressful situations and inflammation. In the first study, greater activity in the dorsal anterior cingulate during a social exclusion task were associated with greater stress-induced increases in the soluble receptor for tumor necrosis factor- α (sTNF-RII), a marker of inflammation, assessed in oral fluids (Slavich, Way, Eisenberger, & Taylor, 2010). In a more recent study, greater amygdala activity during negative social evaluation was related to greater stress-induced increases in plasma IL-6 (Muscatell et al., 2015). These studies together provide initial evidence that the amygdala and PFC are related to stress-related inflammatory processes. Future work should substantiate these findings, in addition to clarifying the roles of early adversity and HPA axis and ANS function in these links.

Early adversity-related alterations in neural function may promote more frequent detection of threat and prolonged and/or exaggerated responses to threat, which can lead to sustained HPA and SNS activation and PNS withdrawal. Early adversity related decreased hippocampal and PFC volume may also interfere with the negative feedback loop of the HPA axis, resulting

in higher levels of cortisol. Over time, these biological states can promote inflammation. Given that sympathetic activity can upregulate inflammatory processes and parasympathetic activity can downregulate inflammatory processes, exaggerated and sustained activation of the SNS and sustained PNS withdrawal can contribute to increases in inflammation. Although cortisol typically has anti-inflammatory effects, prolonged exposure to stress, presumably accompanied by sustained activation of the HPA axis, can desensitize glucocorticoid receptors, which could result in inadequate suppression of inflammation (Cohen et al., 2012).

Although the studies described in the present review collectively suggest that early adversity related changes in the development of the amygdala, hippocampus, and mPFC may mediate the early adversity-inflammation association, no study to date has actually incorporated assessment of all the neurobiological systems; therefore, empirical support for this pathway is currently nonexistent. However, studies are beginning to chart this pathway by using gene expression in immune cells as a window into the dynamics between early adversity-related alterations in neuroendocrine and immune functioning. In one study (Cole et al., 2012), compared to rhesus macaques that were maternally-reared, those that were peer-reared exhibited greater inflammatory gene expression, and greater activity of CREB transcription factors, which play an essential role in the effects of the SNS. Interestingly, there was no evidence of decreased glucocorticoid signaling. Similarly, in a study of human adults (Miller et al., 2009), low socioeconomic status during childhood, which is often associated with a harsh family environment (Dodge, Pettit, & Bates, 1994; Pinderhughes, Dodge, Zelli, Bates, & Pettit, 2000), was also related to greater inflammatory gene expression and CREB signaling. In this study, lower socioeconomic status was also related to decreased glucocorticoid receptor signaling. Further support that increased inflammation resulting from early adversity may be mediated by decreased glucocorticoid receptor sensitivity comes from a longitudinal study of female adolescents raised in either a harsh or nurturing family environment (Miller & Chen, 2010). In addition to circulating levels of pro-inflammatory cytokines, this study examined the capacity of immune cells to mount an inflammatory response to microbial challenge *in vitro* (i.e., stimulated production of pro-inflammatory cytokines). Harsh family climate related changes in circulating levels of inflammation were not observed. Over time, however, the white blood cells of those raised in a harsh family climate exhibited increasingly exaggerated inflammatory responses to bacterial stimuli and greater resistance to the anti-

inflammatory effects of cortisol *in vitro*. This suggests that early adversity may lead immune cells to develop a pro-inflammatory tendency relatively early in life, which may contribute to a chronic pro-inflammatory state later in life (Miller et al., 2011). Collectively, this study and those examining gene expression in relation to early adversity offer initial evidence that early adversity may upregulate SNS activity and desensitize glucocorticoid receptors to increase inflammation and are an important step towards understanding the role of alterations in neural development in the early adversity-inflammation link. Assessing neural structure and function in similar such studies is a critical next step in this line of work.

INFLAMMATORY REGULATION OF NEURAL DEVELOPMENT

The main focus of the present review is to consider the implications of altered neural development related to early adversity for peripheral inflammatory processes. Nevertheless, the bidirectional interactions between neural and inflammatory processes require consideration of the role that inflammatory processes play in neural development.

Inflammatory cytokines and other immune molecules play a critical role in various aspects of neural development, including neural plasticity, neurogenesis, dendrite development, and hippocampal long-term potentiation (Gilmore, Jarskog, Vadlamudi, & Lauder, 2004; Stolp, 2013; Yirmiya & Goshen, 2011). Thus, increased inflammation during early life when both the central nervous and immune systems are developing can have long-lasting detrimental effects on cognitive functioning and behavior and increase risk for developmental neuropsychiatric disorders, including schizophrenia and autism spectrum disorder (Meyer, Feldon, & Dammann, 2011; Yirmiya & Goshen, 2011). For instance, rodent experimental studies show that neonatal and postnatal infection can prime microglia, immune cells located in the brain, to secrete more inflammatory molecules; therefore, when the immune system is subsequently activated, the heightened levels of inflammatory molecules interfere with learning and memory (Bilbo, 2013). Similarly, LPS- and PolyI:C-induced inflammation in the absence of infection in pregnant mothers alters inflammatory cytokines in the fetal brain (Smith, Li, Garbett, Mirnics, & Patterson, 2007; Garay, Hsiao, Patterson, & McAllister, 2013). These changes can produce neuroanatomical, neurochemical, behavioral, and cognitive changes implicated in schizophrenia and autism in offspring. More specifically, they can disrupt dopamine and GABA systems,

social interactions, working memory, selective attention, and sensorimotor gating (Feigenson, Kusnecov, & Silverstein, 2014; Malkova, Yu, Hsiao, Moore, & Patterson, 2012; Meyer & Feldon, 2009, 2012). Whether early life psychological stress-induced inflammation has similar effects is not entirely clear.

In human studies, endotoxin- and typhoid-induced increases in systemic inflammation have been shown to affect neural substrates, particularly those underlying socio-emotional processes. Increased inflammation has been related to increased amygdala and subgenual anterior cingulate cortex activation to socially threatening stimuli (Harrison et al., 2009; Inagaki, Muscatell, Irwin, Cole, & Eisenberger, 2012). Greater amygdala activation to social threat, in turn, was correlated with increased feelings of social disconnection (Inagaki et al., 2012), and greater subgenual anterior cingulate cortex activation was correlated with greater mood deterioration (Harrison et al., 2009). Social disconnection, negative affect, and a lack of positive affect, have also been shown to upregulate inflammation (Chiang et al., 2015; Howren, Lamkin, & Suls, 2009; Jaremka et al., 2013; Steptoe, O'Donnell, Badrick, Kumari, & Marmot, 2008). Whether early adversity influences these associations has not yet been determined.

These findings underscore the fact that inflammation also influences neural development and processes. How increased inflammation following early psychological stress may affect neural development remains incompletely understood. Yet the fact that there are bidirectional effects of the CNS and immune systems makes plausible the notion that early adversity may create a cycle of dysregulation in these neurobiological systems.

FUTURE DIRECTIONS

The present review considered how alterations in neural development associated with early adversity may lead to increased inflammation. Past work has contributed significantly to our understanding of how early adversity might affect inflammatory processes and therefore physical health, but a number of issues remain.

Longitudinal Assessment of Multiple Systems

The majority of studies examining early adversity and neurobiological systems have been cross-sectional. Moreover, to our knowledge, there are no studies to date that have examined changes in neural development as a mechanism linking early adversity and increased inflammation. Thus, longitudinal investigations of early adversity involving repeated assessment of neural structure and functioning and markers of the HPA axis,

ANS, and inflammation are needed. An integrative approach in a prospective design would help elucidate how early stressful experiences affect developmental trajectories of neurobiological systems that ultimately culminate in increased inflammation and greater risk for poor health. It would be relatively easy to add assessment of the HPA axis, SNS, and inflammatory processes to neuroimaging studies that examine the effects of early adversity on neural development. Cortisol and α -amylase, a digestive enzyme secreted by salivary glands that are under SNS control, can be assessed via saliva samples, whereas assessment of inflammatory markers such as CRP can be assessed from dried blood spots (DBS) (McDade, Burhop, & Dohmal, 2004). Both saliva sampling and DBS collection are relatively non-invasive procedures. In fact, DBS requires no medical personnel and has been used routinely to perform medical screening tests in infants. Additionally, assaying saliva samples for cortisol and α -amylase and DBS for CRP is relatively inexpensive.

Neural Connectivity

Studies of the effects of early adversity on connectivity between the amygdala, hippocampus, and mPFC are uncommon, despite the fact that these connections are implicated in contextual learning memory processes pertinent to fear conditioning and extinction (Maren et al., 2013; Stein et al., 2007). Given that memory and fear extinction are believed to play a role in appraisal and reactivity to subsequent contextual stimuli, alterations in connectivity among the amygdala, hippocampus, and PFC and their functional interactions may have implications for HPA and ANS activity during subsequent stress, and ultimately, for inflammatory processes. A research priority for next studies, then, is to investigate the effects of early adversity on amygdala-hippocampal-PFC structural and functional connectivity and their impact on HPA, ANS, and inflammatory processes.

Sensitive Periods

The impact of child maltreatment on neural development may depend on timing of exposure of maltreatment and the temporal development of specific neural regions (Lupien et al., 2009; Payne, Machado, Bliwise, & Bachevalier, 2010). The amygdala, hippocampus, and PFC all undergo significant development during childhood and adolescence, but specific periods of rapid development differ across these regions. The amygdala and hippocampus undergo rapid development earlier in childhood (Tottenham & Sheridan, 2010), whereas the PFC undergoes significant reorganization during adolescence (Blakemore & Choudhury, 2006). As a result,

exposure to stress during these periods may have differential effects on the different neural regions. Indeed, early adversity during late (8–17 years) rather than early (1 month–7 years) childhood has been associated with smaller PFC volume whereas early adversity during early childhood has been associated with smaller hippocampal volume (Andersen et al., 2008; Baker et al., 2012). Determining whether early stressful experiences during particular developmental stages have differential and/or additive effects on neurobiological systems can help determine optimal times to intervene.

Severity of Early Adversity

A particular gap in the literature concerns the effects of normal variation in parenting on neurobiological systems. Past studies have almost exclusively focused on traumatic events and severe forms of maltreatment (i.e., physical, emotional, and sexual abuse and neglect). Yet, most children do not experience severe forms of maltreatment, but may nevertheless experience negative relationships with their parents and other stressors (e.g., Felitti et al., 1998; Repetti et al., 2002). For instance, spanking without use of any objects is a common form of discipline in the United States (Zolotor, Theodore, Runyan, Chang, & Laskey, 2011). Rodent studies suggest that typical variations in parenting (i.e., licking and grooming) can lead to differences in the expression of hippocampal glucocorticoid receptors and the ability to induce synaptic potentiation, thereby affecting HPA and hippocampal functioning (van Hasselt et al., 2012; Weaver et al., 2004). Similarly, in a human study, corporal punishment not considered physical abuse was associated with structural alterations in the anterior cingulate cortex (Tomoda et al., 2009). Thus, further investigation of particular aspects of normative parenting that adversely affect neurobiological systems is warranted.

Genetic Factors

The role that genetics play in these processes and which polymorphisms might contribute to the development of stress-related neurobiological vulnerabilities that may culminate in inflammation are beginning to be elucidated. Single nucleotide polymorphisms in the BDNF and serotonin transporter genes have been found to moderate vulnerability to alterations in developing neurobiological systems as a result of early adversity. BDNF is critical for neuronal growth and survival, supports learning and memory, and plays an important role in inflammatory processes. As a result, the Met allele, which leads to lower levels of BDNF secretion relative to the Val allele, may enhance vulnerability to

the neural and inflammatory effects of early adversity. Indeed, Met carriers with a history of early life stress have smaller hippocampal, amygdala, and anterior cingulate cortical volumes compared to homozygous Val carriers with a history of early life stress (Carballedo et al., 2013; Gatt et al., 2009; Gerritsen et al., 2011). Whether greater levels of inflammation are exhibited among Met carriers compared to Val carriers remains to be seen.

The serotonin transporter gene, which regulates transportation of the neurotransmitter serotonin, may also increase vulnerability to the negative neural effects of early life stress. Depressed S allele carriers who experienced early adversity in the form of emotional neglect were found to have smaller hippocampal and prefrontal cortical volumes compared to those with either a genetic or environmental risk factor (Frodl et al., 2010). As work on genetics progresses, additional genetic polymorphisms that unduly differentiate vulnerability to early family adversity are likely to be identified.

CONCLUSION

Alterations in neural development of the amygdala, hippocampus, and PFC may be a key pathway linking early adversity to not only poor mental health, but also to greater inflammation and subsequent poor physical health. Separate lines of work together support this hypothesis, as described in this review. Specifically, studies have begun to document links between early adversity and alterations in functional and structural development of the amygdala, hippocampus, and PFC, HPA and ANS functioning, and inflammatory processes. Based on these data and on known anatomical connections and physiological regulations among these neurobiological systems, we conclude that altered neural development resulting from early adversity likely ultimately increases inflammation and risk for poor health outcomes in adulthood.

REFERENCES

- Abboud, F. M., Harwani, S. C., & Chapleau, M. W. (2012). Autonomic neural regulation of the immune system: Implications for hypertension and cardiovascular disease. *Hypertension*, 59(4), 755–762.
- Anda, R. F., Felitti, V. J., Bremner, J. D., Walker, J. D., Whitfield, C. H., Perry, B. D., . . . Giles, W. H. (2006). The enduring effects of abuse and related adverse experiences in childhood. *European Archives of Psychiatry and Clinical Neuroscience*, 256(3), 174–186.

- Andersen, S. L., & Teicher, M. H. (2004). Delayed effects of early stress on hippocampal development. *Neuropsychopharmacology*, 29(11), 1988–1993.
- Andersen, S. L., & Teicher, M. H. (2009). Desperately driven and no brakes: developmental stress exposure and subsequent risk for substance abuse. *Neuroscience & Biobehavioral Reviews*, 33(4), 516–524.
- Andersen, S., Tomada, A., Vincow, E., Valente, E., Polcari, A., & Teicher, M. (2008). Preliminary evidence for sensitive periods in the effect of childhood sexual abuse on regional brain development. *The Journal of Neuro-psychiatry and Clinical Neurosciences*, 20(3), 292–301.
- Ashman, S. B., Dawson, G., Panagiotides, H., Yamada, E., & Wilkinson, C. W. (2002). Stress hormone levels of children of depressed mothers. *Development and Psychopathology*, 14(2), 333–349.
- Avitsur, R., Stark, J. L., & Sheridan, J. F. (2001). Social stress induces glucocorticoid resistance in subordinate animals. *Hormones and Behavior*, 39(4), 247–257.
- Baker, L. M., Williams, L. M., Korgaonkar, M. S., Cohen, R. A., Heaps, J. M., & Paul, R. H. (2013). Impact of early vs. late childhood early life stress on brain morphometrics. *Brain Imaging and Behavior*, 7(2), 196–203.
- Bierhaus, A., Wolf, J., Andrassy, M., Rohleder, N., Humpert, P. M., Petrov, D., . . . Nawroth, P. P. (2003). A mechanism converting psychosocial stress into mononuclear cell activation. *Proceedings of the National Academy of Sciences*, 100(4), 1920–1925.
- Bilbo, Staci D. (2013). FRANK A BEACH AWARD: Programming of neuroendocrine function by early-life experience: A critical role for the immune system. *Hormones and Behavior*, 65(5), 684–691.
- Blakemore, S. J., & Choudhury, S. (2006). Development of the adolescent brain: implications for executive function and social cognition. *Journal of Child Psychology and Psychiatry*, 47(3-4), 296–312.
- Bos, K. J., Fox, N., Zeanah, C. H., & Nelson, C. A., III. (2009). Effects of early psychosocial deprivation on the development of memory and executive function. *Frontiers in Behavioral Neuroscience*, 3, 1–7.
- Brake, W. G., Flores, G., Francis, D., Meaney, M. J., Srivastava, L. K., & Gratton, A. (2000). Enhanced nucleus accumbens dopamine and plasma corticosterone stress responses in adult rats with neonatal excitotoxic lesions to the medial prefrontal cortex. *Neuroscience*, 96(4), 687–695.
- Buchanan, T. W., Kern, S., Allen, J. S., Tranel, D., & Kirschbaum, C. (2004). Circadian regulation of cortisol after hippocampal damage in humans. *Biological Psychiatry*, 56(9), 651–656.
- Calkins, S. D., Smith, C. L., Gill, K. L., & Johnson, M. C. (1998). Maternal interactive style across contexts: Relations to emotional, behavioral and physiological regulation during toddlerhood. *Social Development*, 7(3), 350–369.
- Calkins, S. D., Graziano, P. A., Berdan, L. E., Keane, S. P., & Degnan, K. A. (2008). Predicting cardiac vagal regulation in early childhood from maternal–child relationship quality during toddlerhood. *Developmental Psychobiology*, 50(8), 751–766.
- Carballedo, A., Morris, D., Zill, P., Fahey, C., Reinhold, E., Meisenzahl, E., . . . Frodl, T. (2013). Brain-derived neurotrophic factor Val66Met polymorphism and early life adversity affect hippocampal volume. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 162, 183–190.
- Carpenter, L. L., Carvalho, J. P., Tyrka, A. R., Wier, L. M., Mello, A. F., Mello, M. F., . . . Price, L. H. (2007). Decreased adrenocorticotropic hormone and cortisol responses to stress in healthy adults reporting significant childhood maltreatment. *Biological Psychiatry*, 62(10), 1080–1087.
- Carpenter, L. L., Tyrka, A. R., Ross, N. S., Khoury, L., Anderson, G. M., & Price, L. H. (2009). Effect of childhood emotional abuse and age on cortisol responsivity in adulthood. *Biological Psychiatry*, 66(1), 69–75.
- Carpenter, L. L., Gawuga, C. E., Tyrka, A. R., Lee, J. K., Anderson, G. M., & Price, L. H. (2010). Association between plasma IL-6 response to acute stress and early-life adversity in healthy adults. *Neuropsychopharmacology*, 35(13), 2617–2623.
- Carpenter, L. L., Shattuck, T. T., Tyrka, A. R., Geraciotti, T. D., & Price, L. H. (2011). Effect of childhood physical abuse on cortisol stress response. *Psychopharmacology (Berl)*, 214(1), 367–375.
- Carrion, V. G., Weems, C. F., Richert, K., Hoffman, B. C., & Reiss, A. L. (2010). Decreased prefrontal cortical volume associated with increased bedtime cortisol in traumatized youth. *Biological Psychiatry*, 68(5), 491–493.
- Chadwick, M. J., Hassabis, D., Weiskopf, N., & Maguire, E. A. (2010). Decoding individual episodic memory traces in the human hippocampus. *Current Biology*, 20(6), 544–547.
- Chaney, A., Carballedo, A., Amico, F., Fagan, A., Skokauskas, N., Meaney, J., & Frodl, T. (2014). Effect of childhood maltreatment on brain structure in adult patients with major depressive disorder and healthy participants. *Journal of Psychiatry Neuroscience*, 39(1), 50–59.
- Chiang, J. J., Bower, J. E., Almeida, D. M., Irwin, M. R., Seeman, T., & Fuligni, A. J. (2015). Socioeconomic status, daily affective and social experiences, and inflammation during adolescence. *Psychosomatic Medicine*, 77(3), 256–66.
- Chugani, H. T., Behen, M. E., Muzik, O., Juhász, C., Nagy, F., & Chugani, D. C. (2001). Local brain functional activity following early deprivation: a study of post-institutionalized Romanian orphans. *NeuroImage*, 14(6), 1290–1301.
- Cohen, R. A., Grieve, S., Hoth, K. F., Paul, R. H., Sweet, L., Tate, D., . . . Williams, L. M. (2006). Early life stress and morphometry of the adult anterior cingulate cortex and caudate nuclei. *Biological Psychiatry*, 59(10), 975–982. DOI: 10.1016/j.biopsych.2005.12.016
- Cohen, S., Janicki-Deverts, D., Doyle, W. J., Miller, G. E., Frank, E., Rabin, B. S., & Turner, R. B. (2012). Chronic stress, glucocorticoid receptor resistance, inflammation,

- and disease risk. *Proceedings of the National Academy of Sciences*, 109(16), 5995–5999.
- Cole, S. W., Arevalo, J. M. G., Takahashi, R., Sloan, E. K., Lutgendorf, S. K., Sood, A. K., . . . Seeman, T. E. (2010). Computational identification of gene-social environment interaction at the human IL6 locus. *Proceedings of the National Academy of Sciences*, 107(12), 5681–5686.
- Cole, S. W., Conti, G., Arevalo, J. M. G., Ruggiero, A. M., Heckman, J. J., & Suomi, S. J. (2012). Transcriptional modulation of the developing immune system by early life social adversity. *Proceedings of the National Academy of Sciences*, 109(50), 20578–20583.
- Copeland, W. E., Wolke, D., Lereya, S. T., Shanahan, L., Worthman, C. M., & Costello, J. E. (2014). Childhood bullying involvement predicts low-grade systemic inflammation into adulthood. *Proceedings of the National Academy of Sciences*, 111(21), 7570–7575.
- Coussens, L. M., & Werb, Z. (2002). Inflammation and cancer. *Nature*, 420(6917), 860–867.
- Dale, L. P., Carroll, L. E., Galen, G., Hayes, J. A., Webb, K. W., & Porges, S. W. (2009). Abuse history is related to autonomic regulation to mild exercise and psychological wellbeing. *Applied Psychophysiology and Biofeedback*, 34(4), 299–308.
- Danese, A., & McEwen, B. S. (2012). Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiology & Behavior*, 106(1), 29–39.
- Danese, A., Pariante, C. M., Caspi, A., Taylor, A., & Poulton, R. (2007). Childhood maltreatment predicts adult inflammation in a life-course study. *Proceedings of the National Academy of Sciences*, 104(4), 1319–1324. DOI: 10.1073/pnas.0610362104
- Dannlowski, U., Stuhrmann, A., Beutelmann, V., Zwanzger, P., Lenzen, T., Grotegerd, D., & Kugel, H. (2012). Limbic scars: long-term consequences of childhood maltreatment revealed by functional and structural magnetic resonance imaging. *Biological Psychiatry*, 71(4), 286–293.
- Dannlowski, U., Kugel, H., Huber, F., Stuhrmann, A., Redlich, R., Grotegerd, D., . . . Suslow, T. (2013). Childhood maltreatment is associated with an automatic negative emotion processing bias in the amygdala. *Human Brain Mapping*, 34(11), 2899–2909.
- Davies, P. T., & Cicchetti, D. (2004). Toward an integration of family systems and developmental psychopathology approaches. *Development and Psychopathology*, 16, 477–481.
- De Brito, S. A., Viding, E., Sebastian, C. L., Kelly, P. A., Mechelli, A., Maris, H., & McCrory, E. J. (2013). Reduced orbitofrontal and temporal grey matter in a community sample of maltreated children. *Journal of Child Psychology and Psychiatry*, 54(1), 105–112.
- Deverman, B. E., & Patterson, P. H. (2009). Cytokines and CNS Development. *Neuron*, 64(1), 61–78.
- Dietz, L. J., Stoyak, S., Melhem, N., Porta, G., Matthews, K. A., Payne, M. Walker, & Brent, D. A. (2013). Cortisol response to social stress in parentally bereaved youth. *Biological Psychiatry*, 73(4), 379–387.
- Diorio, D., Viau, V., & Meaney, M. J. (1993). The role of the medial prefrontal cortex (cingulate gyrus) in the regulation of hypothalamic-pituitary-adrenal responses to stress. *The Journal of Neuroscience*, 13(9), 3839–3847.
- Dodge, K. A., Pettit, G. S., & Bates, J. E. (1994). Socialization mediators of the relation between socioeconomic status and child conduct problems. *Child Development*, 65(2), 649–655.
- Doom, J. R., Cicchetti, D., & Rogosch, F. A. (2014). Longitudinal patterns of cortisol regulation differ in maltreated and nonmaltreated children. *Journal of the American Academy of Child & Adolescent Psychiatry*, 53(11), 1206–1215.
- Edmiston, E. K., & Blackford, J. U. (2013). Childhood maltreatment and response to novel face stimuli presented during functional magnetic resonance imaging in adults. *Psychiatry Research: Neuroimaging*, 212(1), 36–42.
- Edmiston, E. E., Wang, F., Mazure, C. M., Guiney, J., Sinha, R., Mayes, L. C., & Blumberg, H. P. (2011). Corticostriatal-limbic gray matter morphology in adolescents with self-reported exposure to childhood maltreatment. *Archives of Pediatrics & Adolescent Medicine*, 165(12), 1069–1077.
- El-Sheikh, M. (2005). The role of emotional responses and physiological reactivity in the marital conflict-child functioning link. *Journal of Child Psychology and Psychiatry*, 46(11), 1191–1199.
- Ellis, B. J., Essex, M. J., & Boyce, W. T. (2005). Biological sensitivity to context: II. Empirical explorations of an evolutionary-developmental theory. *Development and Psychopathology*, 17(02), 303–328.
- Engert, V., Buss, C., Khalili-Mahani, N., Wadiwalla, M., Dedovic, K., & Pruessner, J. C. (2010a). Investigating the association between early life parental care and stress responsivity in adulthood. *Developmental Neuropsychology*, 35(5), 570–581.
- Engert, V., Efanov, S. I., Dedovic, K., Duchesne, A., Dagher, A., & Pruessner, J. C. (2010b). Perceived early-life maternal care and the cortisol response to repeated psychosocial stress. *Journal of Psychiatry & Neuroscience*, 35(6), 370.
- Essex, M. J., Klein, M. H., Cho, E., & Kalin, N. H. (2002). Maternal stress beginning in infancy may sensitize children to later stress exposure: effects on cortisol and behavior. *Biological Psychiatry*, 52(8), 776–784.
- Fagundes, C. P., & Way, B. M. (2014). Early-life stress and adult inflammation. *Current Directions in Psychological Science*, 23(4), 277–283.
- Fagundes, C. P., Glaser, R., & Kiecolt-Glaser, J. K. (2013). Stressful early life experiences and immune dysregulation across the lifespan. *Brain, Behavior, and Immunity*, 27(1), 8–12. DOI: 10.1016/j.bbi.2012.06.014
- Fanselow, M. S., & Gale, G. D. (2003). The amygdala, fear, and memory. *Annals of the New York Academy of Sciences*, 985(1), 125–134.
- Feigenson, K. A., Kusnecov, A. W., & Silverstein, S. M. (2014). Inflammation and the two-hit hypothesis of schizophrenia. *Neuroscience & Biobehavioral Reviews*, 38, 72–93.

- Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., ... Marks, J.S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, 14(4), 245–258.
- Fendler, K., Karmos, G., & Telegdy, G. (1961). The effect of hippocampal lesion on pituitary-adrenal function. *Acta Physiologica Academiae Scientiarum Hungaricae*, 20, 293–297.
- Fernald, L. C. H., Burke, H. M., & Gunnar, M. R. (2008). Salivary cortisol levels in children of low-income women with high depressive symptomatology. *Development and Psychopathology*, 20(2), 423.
- Figueiredo, H. F., Bruestle, A., Bodie, B., Dolgas, C. M., & Herman, J. P. (2003). The medial prefrontal cortex differentially regulates stress-induced c-fos expression in the forebrain depending on type of stressor. *European Journal of Neuroscience*, 18(8), 2357–2364.
- Fischette, C. T., Komisaruk, B. R., Edinger, H. M., Feder, H. H., & Siegel, A. (1980). Differential fornix ablations and the circadian rhythmicity of adrenal corticosteroid secretion. *Brain Research*, 195(2), 373–387.
- Fox, S. E., Levitt, P., & Nelson III, C. A. (2010). How the timing and quality of early experiences influence the development of brain architecture. *Child Development*, 81(1), 28–40.
- Frodl, T., & O'Keane, V. (2013). How does the brain deal with cumulative stress? A review with focus on developmental stress, HPA axis function and hippocampal structure in humans. *Neurobiology of Disease*, 52, 24–37.
- Frodl, T., Reinhold, E., Koutsouleris, N., Donohoe, G., Bondy, B., Reiser, M., ... Meisenzahl, E. M. (2010). Childhood stress, serotonin transporter gene and brain structures in major depression. *Neuropsychopharmacology*, 35(6), 1383–1390.
- Frodl, T., Reinhold, E., Koutsouleris, N., Reiser, M., & Meisenzahl, E. M. (2010). Interaction of childhood stress with hippocampus and prefrontal cortex volume reduction in major depression. *Journal of Psychiatric Research*, 44(13), 799–807.
- Fuchs, E., & Flügge, G. (2003). Chronic social stress: effects on limbic brain structures. *Physiology & Behavior*, 79(3), 417–427.
- Gaab, J., Rohleder, N., Nater, U. M., & Ehlert, U. (2005). Psychological determinants of the cortisol stress response: the role of anticipatory cognitive appraisal. *Psychoneuroendocrinology*, 30(6), 599–610.
- Garay, P. A., Hsiao, E. Y., Patterson, P. H., & McAllister, A. K. (2013). Maternal immune activation causes age- and region-specific changes in brain cytokines in offspring throughout development. *Brain, Behavior, & Immunity*, 31, 54–68.
- Gatt, J. M., Nemeroff, C. B., Dobson-Stone, C., Paul, R. H., Bryant, R. A., Schofield, P. R., ... Williams, L. M. (2009). Interactions between BDNF Val66Met polymorphism and early life stress predict brain and arousal pathways to syndromal depression and anxiety. *Molecular Psychiatry*, 14(7), 681–695.
- Gee, D. G., Gabard-Durnam, L. J., Flannery, J., Goff, B., Humphreys, K. L., Telzer, E. H., ... Tottenham, N. (2013). Early developmental emergence of human amygdala-prefrontal connectivity after maternal deprivation. *Proceedings of the National Academy of Sciences*, 110(39), 15638–15643.
- Gerritsen, L., Tendolkar, I., Franke, B., Vasquez, A. A., Kooijman, S., Buitelaar, J., ... Rijpkema, M. (2011). BDNF Val66Met genotype modulates the effect of childhood adversity on subgenual anterior cingulate cortex volume in healthy subjects. *Molecular Psychiatry*, 17(6), 597–603.
- Gianaros, P. J., & Sheu, L. K. (2009). A review of neuroimaging studies of stressor-evoked blood pressure reactivity: Emerging evidence for a brain-body pathway to coronary heart disease risk. *Neuroimage*, 47(3), 922–936.
- Gilmore, J. H., Jarskog, L. F., Vadlamudi, S., & Lauder, J. M. (2004). Prenatal infection and risk for schizophrenia: IL-1, IL-6, and TNF inhibit cortical neuron dendrite development. *Neuropsychopharmacology*, 29, (1221–1229).
- Glezer, I., & Rivest, S. (2004). Glucocorticoids: protectors of the brain during innate immune responses. *The Neuroscientist*, 10(6), 538–552.
- Gouin, J., Glaser, R., Malarkey, W. B., Beversdorf, D., & Kiecolt-Glaser, J. K. (2012). Childhood abuse and inflammatory responses to daily stressors. *Annals of Behavioral Medicine*, 44(2), 287–292.
- Greenough, W. T., Black, J. E., & Wallace, C. S. (1987). Experience and brain development. *Child Development*, 58, 539–559.
- Grisanti, L. A., Woster, A. P., Dahlman, J., Sauter, E. R., Combs, C. K., & Porter, J. E. (2011). α 1-adrenergic receptors positively regulate Toll-like receptor cytokine production from human monocytes and macrophages. *Journal of Pharmacology and Experimental Therapeutics*, 338(2), 648–657.
- Gunnar, M., & Quevedo, K. (2007). The neurobiology of stress and development. *Annual Review of Psychology*, 58, 145–173. DOI: 10.1146/annurev.psych.58.110405.085605
- Gunnar, M. R., Wewerka, S., Frenn, K., Long, J. D., & Griggs, C. (2009). Developmental changes in hypothalamus-pituitary-adrenal activity over the transition to adolescence: normative changes and associations with puberty. *Development and Psychopathology*, 21(1), 69–85.
- Hänsel, A., & von Känel, R. (2008). The ventro-medial prefrontal cortex: a major link between the autonomic nervous system, regulation of emotion, and stress reactivity. *Biopsychosocial Medicine*, 2, 21–25.
- Haensel, A., Mills, P. J., Nelesen, R. A., Ziegler, M. G., & Dimsdale, J. E. (2008). The relationship between heart rate variability and inflammatory markers in cardiovascular diseases. *Psychoneuroendocrinology*, 33(10), 1305–1312.
- Hanahan, D., & Weinberg, R. A. (2011). Hallmarks of cancer: the next generation. *Cell*, 144(5), 646–674.

- Hanson, J. L., Chung, M. K., Avants, B. B., Shurtcliff, E. A., Gee, J. C., Davidson, R. J., & Pollak, S. D. (2010). Early stress is associated with alterations in the orbitofrontal cortex: a tensor-based morphometry investigation of brain structure and behavioral risk. *The Journal of Neuroscience*, 30(22), 7466–7472. DOI: 10.1523/JNEUROSCI.0859-10.2010
- Hanson, J. L., Chung, M. K., Avants, B. B., Rudolph, K. D., Shurtcliff, E. A., Gee, J. C., ... Pollak, S. D. (2012). Structural variations in prefrontal cortex mediate the relationship between early childhood stress and spatial working memory. *The Journal of Neuroscience*, 32(23), 7917–7925.
- Hariri, A. R., Mattay, V. S., Tessitore, A., Fera, F., & Weinberger, D. R. (2003). Neocortical modulation of the amygdala response to fearful stimuli. *Biological Psychiatry*, 53(6), 494–501.
- Harrison, N. A., Brydon, L., Walker, C., Gray, M. A., Steptoe, A., & Critchley, H. D. (2009). Inflammation causes mood changes through alterations in subgenual cingulate activity and mesolimbic connectivity. *Biological Psychiatry*, 66(5), 407–414.
- Harrison, D. G., Guzik, T. J., Lob, H. E., Madhur, M. S., Marvar, P. J., Thabet, S. R., ... Weyand, C. M. (2011). Inflammation, immunity, and hypertension. *Hypertension*, 57, (132–140).
- Hart, H., & Rubia, K. (2012). Neuroimaging of child abuse: a critical review. *Frontiers in Human Neuroscience*, 6(52), 1–24. DOI: 10.3389/fnhum.2012.00052
- Heim, C., Newport, D. J., Heit, S., Graham, Y. P., Wilcox, M., Bonsall, R., ... Nemeroff, C. B. (2000). Pituitary-adrenal and autonomic responses to stress in women after sexual and physical abuse in childhood. *Journal of the American Medical Association*, 284(5), 592–597.
- Heim, C., Newport, D. J., Wagner, D., Wilcox, M. M., Miller, A. H., & Nemeroff, C. B. (2002). The role of early adverse experience and adulthood stress in the prediction of neuroendocrine stress reactivity in women: a multiple regression analysis. *Depression and Anxiety*, 15(3), 117–125.
- Herman, J. P., & Cullinan, W. E. (1997). Neurocircuitry of stress: central control of the hypothalamo-pituitary-adrenocortical axis. *Trends in Neurosciences*, 20(2), 78–84.
- Herman, J. P., & Mueller, N. K. (2006). Role of the ventral subiculum in stress integration. *Behavioural Brain Research*, 174(2), 215–224.
- Herman, J. P., Cullinan, W. E., Morano, M. I., Akil, H., & Watson, S. J. (1995). Contribution of the ventral subiculum to inhibitory regulation of the hypothalamo-pituitary-adrenocortical axis. *Journal of Neuroendocrinology*, 7(6), 475–482.
- Herman, J. P., Figueiredo, H., Mueller, N. K., Ulrich-Lai, Y., Ostrander, M. M., Choi, D. C., & Cullinan, W. E. (2003). Central mechanisms of stress integration: hierarchical circuitry controlling hypothalamo-pituitary-adrenocortical responsiveness. *Frontiers in Neuroendocrinology*, 24(3), 151–180.
- Herringa, R. J., Birn, R. M., Ruttle, P. L., Burghy, C. A., Stodola, D. E., Davidson, R. J., & Essex, M. J. (2013). Childhood maltreatment is associated with altered fear circuitry and increased internalizing symptoms by late adolescence. *Proceedings of the National Academy of Sciences*, 110(47), 19119–24.
- Howren, M. B., Lamkin, D. M., & Suls, J. (2009). Associations of depression with C-reactive protein, IL-1, and IL-6: A meta-analysis. *Psychosomatic Medicine*, 71(2), 171–186.
- Inagaki, T. K., Muscatell, K. A., Irwin, M. R., Cole, S. W., & Eisenberger, N. I. (2012). Inflammation selectively enhances amygdala activity to socially threatening images. *Neuroimage*, 59(4), 3222–3226.
- Irwin, M. R., & Cole, S. W. (2011). Reciprocal regulation of the neural and innate immune systems. *Nature Reviews Immunology*, 11(9), 625–632. DOI: 10.1038/nri3042
- Jaremka, L. M., Fagundes, C. P., Peng, J., Bennett, J. M., Glaser, R., Malarkey, W. B., & Kiecolt-Glaser, J. K. (2013). Loneliness promotes inflammation during acute stress. *Psychological Science*, 24(7), 1089–1097.
- Juster, R. P., McEwen, B. S., & Lupien, S. J. (2010). Allostatic load biomarkers of chronic stress and impact on health and cognition. *Neuroscience & Biobehavioral Reviews*, 35(1), 2–16.
- Katz, L. F. (2007). Domestic violence and vagal reactivity to peer provocation. *Biological Psychology*, 74(2), 154.
- Kelly, P. A., Viding, E., Wallace, G. L., Schaer, M., De Brito, S. A., Robustelli, B., & McCrory, E. J. (2013). Cortical thickness, surface area, and gyrification abnormalities in children exposed to maltreatment: Neural markers of vulnerability?. *Biological Psychiatry*, 74(11), 845–852.
- Kern, S., Oakes, T. R., Stone, C. K., McAuliff, E. M., Kirschbaum, C., & Davidson, R. J. (2008). Glucose metabolic changes in the prefrontal cortex are associated with HPA axis response to a psychosocial stressor. *Psychoneuroendocrinology*, 33(4), 517–529.
- Kiecolt-Glaser, J. K., Gouin, J. P., Weng, N., Malarkey, W. B., Beversdorf, D. Q., & Glaser, R. (2011). Childhood adversity heightens the impact of later-life caregiving stress on telomere length and inflammation. *Psychosomatic Medicine*, 73(1), 16–22.
- Kitayama, N., Quinn, S., & Bremner, J. D. (2006). Smaller volume of anterior cingulate cortex in abuse-related posttraumatic stress disorder. *Journal of Affective Disorders*, 90(2), 171–174.
- Knigge, K. M. (1961). Adrenocortical response to stress in rats with lesions in hippocampus and amygdala. *Proceedings of the Society for Experimental Biology and Medicine*, 108, 18–21.
- Korgaonkar, M. S., Antees, C., Williams, L. M., Gatt, J. M., Bryant, R. A., Cohen, R., ... Grieve, S. M. (2013). Early exposure to traumatic stressors impairs emotional brain circuitry. *Public Library of Science One*, 8(9), 1–8.
- Kuhlman, K. R., Geiss, E. G., Vargas, I., & Lopez-Duran, N. L. (2015). Differential associations between childhood trauma subtypes and adolescent HPA-axis functioning. *Psychoneuroendocrinology*, 54, 103–114.

- Kumari, M., Head, J., Bartley, M., Stansfeld, S., & Kivimaki, M. (2012). Maternal separation in childhood and diurnal cortisol patterns in mid-life: findings from the Whitehall II study. *Psychological Medicine*, 1(1), 1–11.
- Kunz-Ebrecht, S. R., Mohamed-Ali, V., Feldman, P. J., Kirschbaum, C., & Steptoe, A. (2003). Cortisol responses to mild psychological stress are inversely associated with proinflammatory cytokines. *Brain, Behavior, and Immunity*, 17(5), 373–383.
- Lacey, R. E., Kumari, M., & Bartley, M. (2014). Social isolation in childhood and adult inflammation: evidence from the National Child Development Study. *Psychoneuroendocrinology*, 50, 85–94.
- Lane, R. D., McRae, K., Reiman, Eric M., Chen, K., Ahern, Geoffrey L., & Thayer, J. F. (2009). Neural correlates of heart rate variability during emotion. *Neuroimage*, 44(1), 213–222.
- Larkin, K. T., Frazer, N. L., & Semenchuk, E. M. (1996). Physiological, affective, and behavioral responses to interpersonal conflict among males from families with different levels of cohesion and adaptability. *Journal of Psychopathology and Behavioral Assessment*, 18(3), 239–254.
- Larkin, K. T., Frazer, N. L., & Wheat, A. L. (2011). Responses to interpersonal conflict among young adults: Influence of family of origin. *Personal Relationships*, 18(4), 657–667.
- LeDoux, J. (2003). The emotional brain, fear, and the amygdala. *Cellular and Molecular Neurobiology*, 23(4–5), 727–738.
- Libby, P. (2006). Inflammation and cardiovascular disease mechanisms. *The American Journal of Clinical Nutrition*, 83(2), 456S–460S.
- Loman, M. M., & Gunnar, M. R. (2010). Early experience and the development of stress reactivity and regulation in children. *Neuroscience & Biobehavioral Reviews*, 34(6), 867–876. DOI: 10.1016/j.neubiorev.2009.05.007.
- Lovallo, W. R., Farag, N. H., Sorocco, K. H., Cohoon, A. J., & Vincent, A. S. (2012). Lifetime adversity leads to blunted stress axis reactivity: studies from the Oklahoma Family Health Patterns Project. *Biological Psychiatry*, 71(4), 344–349.
- Lovallo, W. R. (2013). Early life adversity reduces stress reactivity and enhances impulsive behavior: Implications for health behaviors. *International Journal of Psychophysiology*, 90(1), 8–16.
- Lovejoy, M. C., Graczyk, P. A., O'Hare, E., & Neuman, G. (2000). Maternal depression and parenting behavior: A meta-analytic review. *Clinical Psychology Review*, 20(5), 561–592.
- Luecken, L. J., & Roubinov, D. S. (2012). Hostile behavior links negative childhood family relationships to heart rate reactivity and recovery in young adulthood. *International Journal of Psychophysiology*, 84(2), 172–179.
- Luecken, L. J., Kraft, A., & Hagan, M. J. (2009). Negative relationships in the family-of-origin predict attenuated cortisol in emerging adults. *Hormones and Behavior*, 55(3), 412–417.
- Luecken, L. J. (1998). Childhood attachment and loss experiences affect adult cardiovascular and cortisol function. *Psychosomatic Medicine*, 60(6), 765–772.
- Lupien, S. J., McEwen, B. S., Gunnar, M. R., & Heim, C. (2009). Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature Reviews Neuroscience*, 10(6), 434–445. DOI: 10.1038/nrn2639
- Lupien, S. J., Parent, S., Evans, A. C., Tremblay, R. E., Zelazo, P. D., Corbo, V., ... Séguin, J. (2011). Larger amygdala but no change in hippocampal volume in 10-year-old children exposed to maternal depressive symptomatology since birth. *Proceedings of the National Academy of Sciences*, 108(34), 14324–14329.
- MacLulich, A. M. J., Ferguson, K. J., Wardlaw, J. M., Starr, J. M., Deary, I. J., & Seckl, J. R. (2006). Smaller left anterior cingulate cortex volumes are associated with impaired hypothalamic-pituitary-adrenal axis regulation in healthy elderly men. *Journal of Clinical Endocrinology & Metabolism*, 91(4), 1591–1594.
- MacMillan, H. L., Georgiades, K., Duku, E. K., Shea, A., Steiner, M., Niec, A., ... Schmidt, L. A. (2009). Cortisol response to stress in female youths exposed to childhood maltreatment: results of the youth mood project. *Biological Psychiatry*, 66(1), 62–68.
- Magarinos, A. M., Somoza, G., & De Nicola, A. F. (1987). Glucocorticoid negative feedback and glucocorticoid receptors after hippocampectomy in rats. *Hormone and Metabolic Research*, 19(3), 105–109.
- Maheu, F. S., Dozier, M., Guyer, A. E., Mandell, D., Peloso, E., Poeth, K., ... Ernst, M. (2010). A preliminary study of medial temporal lobe function in youths with a history of caregiver deprivation and emotional neglect. *Cognitive, Affective, & Behavioral Neuroscience*, 10(1), 34–49.
- Malkova, N. V., Yu, C. Z., Hsiao, E. Y., Moore, M. J., & Patterson, P. H. (2012). Maternal immune activation yields offspring displaying mouse versions of the three core symptoms of autism. *Brain, Behavior, & Immunity*, 26(4), 607–616.
- Maren, S., Phan, K. L., & Liberzon, I. (2013). The contextual brain: implications for fear conditioning, extinction and psychopathology. *Nature Reviews Neuroscience*, 14(6), 417–428.
- Matthews, K. A., Chang, Y. F., Thurston, R. C., & Bromberger, J. T. (2014). Child abuse is related to inflammation in mid-life women: Role of obesity. *Brain, Behavior, & Immunity*, 36, (29–34).
- McCrory, E., De Brito, S. A., & Viding, E. (2011). The impact of childhood maltreatment: a review of neurobiological and genetic factors. *Frontiers in Psychiatry*, 2, 1–14. DOI: 10.3389/fpsy.2011.00048
- McCrory, E., De Brito, S. A., & Viding, E. (2012). The link between child abuse and psychopathology: a review of neurobiological and genetic research. *Journal of the Royal Society of Medicine*, 105(4), 151–156.
- McDade, T. W., Burhop, J., & Dohnal, J. (2004). High-sensitivity enzyme immunoassay for C-reactive protein in dried blood spots. *Clinical Chemistry*, 50(3), 652–654.

- McEwen, B. S. (1998). Stress, adaptation, and disease: Allostasis and allostatic load. *Annals of the New York Academy of Sciences*, 840(1), 33–44.
- McNamee, E. N., Griffin, E. W., Ryan, K. M., Ryan, K. J., Heffernan, S., Harkin, A., & Connor, T. J. (2010). Noradrenaline acting at β -adrenoceptors induces expression of IL-1 β and its negative regulators IL-1ra and IL-1RII, and drives an overall anti-inflammatory phenotype in rat cortex. *Neuropharmacology*, 59(1-2), 37–48.
- Mehta, M. A., Golemboski, N. I., Nosarti, C., Colvert, E., Mota, A., Williams, S. C. R., . . . Sonuga-Barke, E. J. S. (2009). Amygdala, hippocampal and corpus callosum size following severe early institutional deprivation: the English and Romanian Adoptees study pilot. *Journal of Child Psychology and Psychiatry*, 50(8), 943–951.
- Meyer, U., & Feldon, J. (2009). Prenatal exposure to infection: a primary mechanism for abnormal dopaminergic development in schizophrenia. *Psychopharmacology (Berl)*, 206(4), 587–602.
- Meyer, U., & Feldon, J. (2012). To poly(I:C) or not to poly(I:C): advancing preclinical schizophrenia research through the use of prenatal immune activation models. *Neuropharmacology*, 62(3), 1308–1321.
- Meyer, U., Feldon, J., & Dammann, O. (2011). Schizophrenia and autism: both shared and disorder-specific pathogenesis via perinatal inflammation? *Pediatric Research*, 69(5 Pt 2), 26R–33R.
- Milad, M. R., Wright, C. I., Orr, S. P., Pitman, R. K., Quirk, G. J., & Rauch, S. L. (2007). Recall of fear extinction in humans activates the ventromedial prefrontal cortex and hippocampus in concert. *Biological Psychiatry*, 62(5), 446–454.
- Miller, G. E., & Chen, E. (2010). Harsh family climate in early life presages the emergence of a proinflammatory phenotype in adolescence. *Psychological Science*, 21(6), 848–856.
- Miller, G. E., Cohen, S., & Ritchey, A. K. (2002). Chronic psychological stress and the regulation of pro-inflammatory cytokines: a glucocorticoid-resistance model. *Health Psychology*, 21(6), 531–541.
- Miller, G. E., Chen, E., Fok, A. K., Walker, H., Lim, A., Nicholls, E. F., . . . Kobor, M. S. (2009). Low early-life social class leaves a biological residue manifested by decreased glucocorticoid and increased proinflammatory signaling. *Proceedings of the National Academy of Sciences*, 106(34), 14716–14721.
- Miller, G. E., Chen, E., & Parker, K. J. (2011). Psychological stress in childhood and susceptibility to the chronic diseases of aging: moving toward a model of behavioral and biological mechanisms. *Psychological Bulletin*, 137(6), 959–997. DOI: 10.1037/a0024768
- Miskovic, V., Schmidt, L. A., Georgiades, K., Boyle, M., & MacMillan, H. L. (2009). Stability of resting frontal electroencephalogram (EEG) asymmetry and cardiac vagal tone in adolescent females exposed to child maltreatment. *Developmental Psychobiology*, 51(6), 474–487.
- Moore, G. A. (2010). Parent conflict predicts infants' vagal regulation in social interaction. *Development and Psychopathology*, 22(1), 23–33.
- Moriceau, S., Roth, T. L., & Sullivan, R. M. (2010). Rodent model of infant attachment learning and stress. *Developmental Psychobiology*, 52(7), 651–660.
- Mueller, A., Strahler, J., Armbruster, D., Lesch, K. P., Brocke, B., & Kirschbaum, C. (2012). Genetic contributions to acute autonomic stress responsiveness in children. *International Journal of Psychophysiology*, 83(3), 302–308.
- Musaelyan, K., Egeland, M., Fernandes, C., Pariante, C. M., Zunszain, P. A., & Thuret, S. (2014). Modulation of adult hippocampal neurogenesis by early-life environmental challenges triggering immune activation. *Neural Plasticity*, 1–10.
- Muscattell, K. A., & Eisenberger, N. I. (2012). A social neuroscience perspective on stress and health. *Social and Personality Psychology Compass*, 6(12), 890–904.
- Muscattell, K. A., Dedovic, K., Slavich, G. M., Jarcho, M. R., Breen, E. C., Bower, J. E., . . . Eisenberger, N. I. (2015). Greater amygdala activity and dorsomedial prefrontal-amygdala coupling are associated with enhanced inflammatory responses to stress. *Brain, Behavior, & Immunity*, 43, 46–53.
- Navalta, C. P., Polcari, A., Webster, D. M., Boghossian, A., & Teicher, M. H. (2006). Effects of childhood sexual abuse on neuropsychological and cognitive function in college women. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 18(1), 45–53.
- Oosterman, M., De Schipper, J. C., Fisher, P., Dozier, M., & Schuengel, C. (2010). Autonomic reactivity in relation to attachment and early adversity among foster children. *Development and Psychopathology*, 22(1), 109–118.
- Orsini, C. A., Kim, J. H., Knapska, E., & Maren, S. (2011). Hippocampal and prefrontal projections to the basal amygdala mediate contextual regulation of fear after extinction. *The Journal of Neuroscience*, 31(47), 17269–17277.
- Ouellet-Morin, I., Odgers, C. L., Danese, A., Bowes, L., Shakoor, S., Papadopoulos, A. S., . . . Arseneault, L. (2011). Blunted cortisol responses to stress signal social and behavioral problems among maltreated/bullied 12-year-old children. *Biological Psychiatry*, 70(11), 1016–1023.
- Pace, T., Mletzko, T., Alagbe, O., Musselman, D., Nemeroff, C. B., Miller, A., & Heim, C. (2006). Increased stress-induced inflammatory responses in male patients with major depression and increased early life stress. *American Journal of Psychiatry*, 163(9), 1630–1633.
- Pascual-Leone, A., Amedi, A., Fregni, F., & Merabet, L. B. (2005). The plastic human brain cortex. *Annual Review of Neuroscience*, 28, 377–401.
- Patterson, T. K., Craske, M. G., & Knowlton, B. J. (2013). The effect of early-life stress on memory systems supporting instrumental behavior. *Hippocampus*, 23(11), 1025–1034.
- Payne, C., Machado, C. J., Bliwise, N. G., & Bachevalier, J. (2010). Maturation of the hippocampal formation and amygdala in *Macaca mulatta*: a volumetric magnetic resonance imaging study. *Hippocampus*, 20(8), 922–935.

- Pearson, T. A., Mensah, G. A., Alexander, R. W., Anderson, J. L., Cannon, R. O., Criqui, M., . . . Vinicor, F. (2003). Markers of inflammation and cardiovascular disease application to clinical and public health practice: a statement for healthcare professionals from the centers for disease control and prevention and the American Heart Association. *Circulation*, *107*(3), 499–511.
- Pechtel, P., & Pizzagalli, D. A. (2011). Effects of early life stress on cognitive and affective function: an integrated review of human literature. *Psychopharmacology (Berl)*, *214*(1), 55–70. DOI: 10.1007/s00213-010-2009-2
- Phelps, E. A. (2004). Human emotion and memory: interactions of the amygdala and hippocampal complex. *Current Opinion in Neurobiology*, *14*(2), 198–202.
- Pinderhughes, E. E., Dodge, K. A., Zelli, A., Bates, J. E., & Pettit, G. S. (2000). Discipline responses: Influences of parents' socioeconomic status, ethnicity, beliefs about parenting, stress, and cognitive-emotional processes. *Journal of Family Psychology*, *14*(3), 380–400.
- Pollak, S. D., & Kistler, D. J. (2002). Early experience is associated with the development of categorical representations for facial expressions of emotion. *Proceedings of the National Academy of Sciences*, *99*(13), 9072–9076.
- Porter, C. L., Wouden-Miller, M., Silva, S. S., & Porter, A. E. (2003). Marital harmony and conflict: Links to infants' emotional regulation and cardiac vagal tone. *Infancy*, *4*(2), 297–307.
- Pruessner, J. C., Dedovic, K., Khalili-Mahani, N., Engert, V., Pruessner, M., Buss, C., . . . Lupien, S. (2008). Deactivation of the limbic system during acute psychosocial stress: evidence from positron emission tomography and functional magnetic resonance imaging studies. *Biological Psychiatry*, *63*(2), 234–240.
- Pruessner, J. C., Dedovic, K., Pruessner, M., Lord, C., Buss, C., Collins, L., . . . Lupien, S. J. (2010). Stress regulation in the central nervous system: evidence from structural and functional neuroimaging studies in human populations-2008 Curt Richter Award Winner. *Psychoneuroendocrinology*, *35*(1), 179–191.
- Quirk, G. J., Likhtik, E., Pelletier, J. G., & Paré, D. (2003). Stimulation of medial prefrontal cortex decreases the responsiveness of central amygdala output neurons. *The Journal of Neuroscience*, *23*(25), 8800–8807.
- Radley, J. J., Arias, C. M., & Sawchenko, P. E. (2006). Regional differentiation of the medial prefrontal cortex in regulating adaptive responses to acute emotional stress. *The Journal of Neuroscience*, *26*(50), 12967–12976.
- Raison, C. L., Capuron, L., & Miller, A. H. (2006). Cytokines sing the blues: inflammation and the pathogenesis of depression. *Trends in Immunology*, *27*(1), 24–31.
- Rakoff-Nahoum, S. (2006). Why cancer and inflammation?. *The Yale Journal of Biology and Medicine*, *79*(3-4), 123–130.
- Raposa, E. B., Hammen, C. L., Brennan, P. A., O'Callaghan, F., & Najman, J. M. (2014). Early adversity and health outcomes in young adulthood: The role of ongoing stress. *Health Psychology*, *33*(5), 410–418.
- Repetti, R. L., Taylor, S. E., & Seeman, T. E. (2002). Risky families: Family social environments and the mental and physical health of offspring. *Psychological Bulletin*, *128*(2), 330–366. DOI: 10.1177/074355489272005
- Repetti, R. L., Robles, T. F., & Reynolds, B. (2011). Allostatic processes in the family. *Development and Psychopathology*, *22*, 921–938.
- Ridker, P. M. (2003). Clinical application of C-reactive protein for cardiovascular disease detection and prevention. *Circulation*, *107*(3), 363–369. DOI: 10.1161/01.cir.0000053730.47739.3c
- Rohleder, N. (2011). Variability in stress system regulatory control of inflammation: a critical factor mediating health effects of stress. *Expert Review of Endocrinology & Metabolism*, *6*(2), 269–278.
- Rohleder, N. (2014). Stimulation of systemic low-grade inflammation by psychosocial stress. *Psychosomatic Medicine*, *76*(3), 181–189.
- Romeo, R. D. (2010). Pubertal maturation and programming of hypothalamic–pituitary–adrenal reactivity. *Frontiers in Neuroendocrinology*, *31*(2), 232–240.
- Roosendaal, B., McEwen, B. S., & Chattarji, S. (2009). Stress, memory and the amygdala. *Nature Reviews Neuroscience*, *10*(6), 423–433.
- Rudy, J. W. (1993). Contextual conditioning and auditory cue conditioning dissociate during development. *Behavioral Neuroscience*, *107*(5), 887–891.
- Sapolsky, R. M., Armanini, M. P., Sutton, S. W., & Plotsky, P. M. (1989). Elevation of hypophysial portal concentrations of adrenocorticotropin secretagogues after fornix transection. *Endocrinology*, *125*(6), 2881–2887.
- Schrepf, A., Markon, K., & Lutgendorf, S. K. (2014). From childhood trauma to elevated C-reactive protein in adulthood: The role of anxiety and emotional eating. *Psychosomatic Medicine*, *76*(5), 327–336.
- Shenk, C. E., Noll, J. G., Putnam, F. W., & Trickett, P. K. (2010). A prospective examination of the role of childhood sexual abuse and physiological asymmetry in the development of psychopathology. *Child Abuse & Neglect*, *34*(10), 752–761.
- Sheridan, J. F., Stark, J. L., Avitsur, R., & Padgett, D. A. (2000). Social disruption, immunity, and susceptibility to viral infection: role of glucocorticoid insensitivity and NGF. *Annals of the New York Academy of Sciences*, *917*(1), 894–905.
- Slavich, G. M., Way, B. M., Eisenberger, N. I., & Taylor, S. E. (2010). Neural sensitivity to social rejection is associated with inflammatory responses to social stress. *Proceedings of the National Academy of Sciences*, *107*(33), 14817–14822.
- Slopen, N., Kubzansky, L. D., McLaughlin, K. A., & Koenen, K. C. (2013). Childhood adversity and inflammatory processes in youth: A prospective study. *Psychoneuroendocrinology*, *38*(2), 188–200.
- Smith, A. P., Stephan, K. E., Rugg, M. D., & Dolan, R. J. (2006). Task and content modulate amygdala-hippocampal connectivity in emotional retrieval. *Neuron*, *16*(49), 631–638.

- Smith, S. E. P., Li, J., Garbett, K., Mirmics, K., & Patterson, P. H. (2007). Maternal immune activation alters fetal brain development through interleukin-6. *The Journal of Neuroscience*, 27(40), 10695–10702.
- Sotres-Bayon, F., Cain, C. K., & LeDoux, J. E. (2006). Brain mechanisms of fear extinction: historical perspectives on the contribution of prefrontal cortex. *Biological Psychiatry*, 60(4), 329–336.
- Squire, L. R. (1992). Memory and the hippocampus: a synthesis from findings with rats, monkeys, and humans. *Psychological Review*, 99(2), 195–231.
- Stein, J. L., Wiedholz, L. M., Bassett, D. S., Weinberger, D. R., Zink, C. F., Mattay, V. S., & Meyer-Lindenberg, A. (2007). A validated network of effective amygdala connectivity. *Neuroimage*, 36(3), 736–745.
- Steptoe, A., O'Donnell, K., Badrick, E., Kumari, M., & Marmot, M. (2008). Neuroendocrine and inflammatory factors associated with positive affect in healthy men and women the Whitehall II Study. *American Journal of Epidemiology*, 167(1), 96–102.
- Sternberg, E. M., Chrousos, G. P., Wilder, R. L., & Gold, P. W. (1992). The stress response and the regulation of inflammatory disease. *Annals of Internal Medicine*, 117(10), 854–866.
- Stolp, H. B. (2013). Neurotrophic cytokines in normal brain development and neurodevelopmental disorders. *Molecular and Cellular Neuroscience*, 53, 63–68.
- Stroud, L. R., Foster, E., Papandonatos, G. D., Handwerker, K., Granger, D. A., Kivlighan, K. T., & Niaura, R. (2009). Stress response and the adolescent transition: Performance versus peer rejection stressors. *Development and Psychopathology*, 21(1), 47–68.
- Struber, N., Struber, D., & Rother, G. (2014). Impact of early adversity on glucocorticoid regulation and later mental disorders. *Neuroscience and Biobehavioral Reviews*, 38, 17–37.
- Sullivan, R. M., & Gratton, A. (2002). Prefrontal cortical regulation of hypothalamic–pituitary–adrenal function in the rat and implications for psychopathology: side matters. *Psychoneuroendocrinology*, 27(1), 99–114.
- Sullivan, M. W., Bennett, D. S., & Lewis, M. (2013). Individual differences in the cortisol responses of neglected and comparison children. *Child Maltreatment*, 18(1), 8–16.
- Taylor, S. E., Lerner, J. S., Sage, R. M., Lehman, B. J., & Seeman, T. E. (2004). Early environment, emotions, responses to stress, and health. *Journal of Personality*, 72(6), 1365–1394.
- Taylor, S. E., Eisenberger, N. I., Saxbe, D., Lehman, B. J., & Lieberman, M. D. (2006). Neural responses to emotional stimuli are associated with childhood family stress. *Biological Psychiatry*, 60(3), 296–301. DOI: 10.1016/j.biopsych.2005.09.027
- Taylor, S. E., Lehman, B. J., Kiefe, C. I., & Seeman, T. E. (2006). Relationship of early life stress and psychological functioning to adult C-reactive protein in the coronary artery risk development in young adults study. *Biological Psychiatry*, 60(8), 819–824.
- Taylor, S. E., Karlamangla, A. S., Friedman, E. M., & Seeman, T. E. (2011). Early environment affects neuroendocrine regulation in adulthood. *Social Cognitive and Affective Neuroscience*, 6(2), 244–251.
- Teicher, M. H., Anderson, C. M., & Polcari, A. (2012). Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proceedings of the National Academy of Sciences*, 109(9), E563–E572.
- Thayer, J. F., & Fischer, J. E. (2009). Heart rate variability, overnight urinary norepinephrine and C-reactive protein: evidence for the cholinergic anti-inflammatory pathway in healthy human adults. *Journal of Internal Medicine*, 265(4), 439–447.
- Thayer, J. F., Yamamoto, S. S., & Brosschot, J. F. (2010). The relationship of autonomic imbalance, heart rate variability and cardiovascular disease risk factors. *International Journal of Cardiology*, 141(2), 122–131.
- Thayer, J. F., Åhs, F., Fredrikson, M., Sollers, J. J., III, & Wager, T. D. (2012). A meta-analysis of heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker of stress and health. *Neuroscience & Biobehavioral Reviews*, 36(2), 747–756.
- Thayer, J. F. (2009). Vagal tone and the inflammatory reflex. *Cleveland Clinic Journal of Medicine*, 76(Suppl 2), S23–S26.
- Tomoda, A., Suzuki, H., Rabi, K., Sheu, Y., Polcari, A., & Teicher, M. H. (2009). Reduced prefrontal cortical gray matter volume in young adults exposed to harsh corporal punishment. *Neuroimage*, 47, T66–T71.
- Tottenham, N., & Sheridan, M. A. (2010). A review of adversity, the amygdala and the hippocampus: a consideration of developmental timing. *Frontiers in Human Neuroscience*, 3(68), 1–18. DOI: 10.3389/neuro.09.068.2009
- Tottenham, N., Hare, T. A., Quinn, B. T., McCarry, T. W., Nurse, M., Gilhooly, T., . . . Casey, B. J. (2010). Prolonged institutional rearing is associated with atypically large amygdala volume and difficulties in emotion regulation. *Developmental Science*, 13(1), 46–61.
- Tottenham, N., Hare, T. A., Millner, A., Gilhooly, T., Zevin, J. D., & Casey, B. J. (2011). Elevated amygdala response to faces following early deprivation. *Developmental Science*, 14(2), 190–204. DOI: 10.1111/j.1467-7687.2010.00971.x
- Tracey, K. J. (2002). The inflammatory reflex. *Nature*, 420(6917), 853–859.
- Tracey, K. J. (2009). Reflex control of immunity. *Nature Reviews Immunology*, 9(6), 418–428.
- Treadway, M. T., Grant, M. M., Ding, Z., Hollon, S. D., Gore, J. C., & Shelton, R. C. (2009). Early adverse events, HPA activity and rostral anterior cingulate volume in MDD. *Public Library of Science One*, 4(3), e4887.
- Trickett, P. K., Gordis, E., Peckins, M. K., & Susman, E. J. (2014). Stress reactivity in maltreated and comparison male and female young adolescents. *Child Maltreatment*, 19, 27–37.

- Turney, K. (2011). Labored love: Examining the link between maternal depression and parenting behaviors. *Social Science Research*, 40(1), 399–415.
- Tyrka, A. R., Burgers, D. E., Philip, N. S., Price, L. H., & Carpenter, L. L. (2013). The neurobiological correlates of childhood adversity and implications for treatment. *Acta Psychiatrica Scandinavica*, 128(6), 434–447.
- Urry, H. L., Van Reekum, C. M., Johnstone, T., Kalin, N. H., Thurow, M. E., Schaefer, H. S., . . . Davidson, R. J. (2006). Amygdala and ventromedial prefrontal cortex are inversely coupled during regulation of negative affect and predict the diurnal pattern of cortisol secretion among older adults. *The Journal of Neuroscience*, 26(16), 4415–4425.
- Verberne, A. J. M., & Owens, N. C. (1998). Cortical Modulation of the Cardiovascular System. *Progress in Neurobiology*, 54(2), 149–168.
- van Harmelen, A. L., van Tol, M. J., van der Wee, N. J., Veltman, D. J., Aleman, A., Spinhoven, P., . . . Elzinga, B. M. (2010). Reduced medial prefrontal cortex volume in adults reporting childhood emotional maltreatment. *Biological Psychiatry*, 68(9), 832–838. DOI: 10.1016/j.biopsych.2010.06.011.
- van Harmelen, A. L., van Tol, M. J., Demenescu, L. R., van der Wee, N. J., Veltman, D. J., Aleman, A., & Elzinga, B. M. (2012). Enhanced amygdala reactivity to emotional faces in adults reporting childhood emotional maltreatment. *Social Cognitive and Affective Neuroscience*, DOI: 10.1093/scan/nss007
- van Hasselt, F. N., Cornelisse, S., Yuan Zhang, T., Meaney, M. J., Velzing, E. H., Krugers, H. J., & Joëls, M. (2012). Adult hippocampal glucocorticoid receptor expression and dentate synaptic plasticity correlate with maternal care received by individuals early in life. *Hippocampus*, 22(2), 255–266.
- van der Vegt, E. J. M., van der Ende, J., Kirschbaum, C., Verhulst, F. C., & Tiemeier, H. (2009). Early neglect and abuse predict diurnal cortisol patterns in adults. A study of international adoptees. *Psychoneuroendocrinology*, 34, 660–669.
- Wager, T. D., van Ast, V. A., Hughes, B. L., Davidson, M. L., Lindquist, M. A., & Ochsner, K. N. (2009). Brain mediators of cardiovascular responses to social threat, part II: Prefrontal-subcortical pathways and relationship with anxiety. *Neuroimage*, 47(3), 836–851.
- Wang, J., Rao, H., Wetmore, G. S., Furlan, P. M., Korczykowski, M., Dinges, D. F., & Detre, J. A. (2005). Perfusion functional MRI reveals cerebral blood flow pattern under psychological stress. *Proceedings of the National Academy of Sciences*, 102(49), 17804–17809.
- Weaver, I. C. G., Cervoni, N., Champagne, F. A., D'Alessio, A. C., Sharma, S., Seckl, J. R., . . . Meaney, M. J. (2004). Epigenetic programming by maternal behavior. *Nature Neuroscience*, 7(8), 847–854.
- Weber, C. S., Thayer, J. F., Rudat, M., Wirtz, P. H., Zimmermann-Viehoff, F., Thomas, A., & Deter, H. C. (2010). Low vagal tone is associated with impaired post stress recovery of cardiovascular, endocrine, and immune markers. *European Journal of Applied Physiology*, 109(2), 201–211.
- Wellen, K. E., & Hotamisligil, G. S. (2005). Inflammation, stress, and diabetes. *Journal of Clinical Investigation*, 115(5), 1111–1119.
- Whittle, S., Yap, M. B. H., Yücel, M., Sheeber, L., Simmons, J. G., Pantelis, C., & Allen, N. B. (2009). Maternal responses to adolescent positive affect are associated with adolescents' reward neuroanatomy. *Social Cognitive and Affective Neuroscience*, 4(3), 247–256.
- Yang, J. H., Lee, E. O., Kim, S. E., Suh, Y. H., & Chong, Y. H. (2012). Norepinephrine differentially modulates the innate inflammatory response provoked by amyloid- β peptide via action at β -adrenoceptors and activation of cAMP/PKA pathway in human THP-1 macrophages. *Experimental Neurology*, 236(2), 199–206.
- Yirmiya, R., & Goshen, I. (2011). Immune modulation of learning, memory, neural plasticity and neurogenesis. *Brain, Behavior, and Immunity*, 25(2), 181–213.
- Zagon, A., & Smith, A. D. (1993). Monosynaptic projections from the rostral ventrolateral medulla oblongata to identified sympathetic preganglionic neurons. *Neuroscience*, 54(3), 729–743.
- Zolotor, A. J., Theodore, A. D., Runyan, D. K., Chang, J. J., & Laskey, A. L. (2011). Corporal punishment and physical abuse: population-based trends for three-to-11-year-old children in the United States. *Child Abuse Review*, 20(1), 57–66.